arXiv:1507.05159v2 [math.QA] 31 Jul 2015

An S_3-symmetry of the Jacobi Identity for Intertwining Operator Algebras

Ling Chen

Abstract

We prove an S_3-symmetry of the Jacobi identity for intertwining operator algebras. Since this Jacobi identity involves the braiding and fusing isomorphisms satisfying the genus-zero Moore-Seiberg equations, our proof uses not only the basic properties of intertwining operators, but also the properties of braiding and fusing isomorphisms and the genus-zero Moore-Seiberg equations. Our proof depends heavily on the theory of multivalued analytic functions of several variables, especially the theory of analytic extensions.

Mathematics Subject Classification (2010). 17B69, 81T40.
Key words. Intertwining operator algebras, Moore-Seiberg equations, Jacobi identity, S_3-symmetry.

1 Introduction

Intertwining operator algebras were introduced and studied by Huang in [H1, H2]. In [C], the author studied intertwining operator algebras in a setting more general than [H2]. In particular, the duality properties, Jacobi identity, Moore-Seiberg equations, locality and some other properties of intertwining operator algebras were studied. For the background on intertwining operator algebras, we refer the reader to [H1, H2, C].

For vertex operator algebras, the Jacobi identity has an S_3-symmetry which corresponds to the obvious S_3-symmetry of the Jacobi identity for Lie algebras [FHL]. For abelian intertwining operator algebras (see [DL1, DL2]), Guo [G] proved that the Jacobi identity for these algebras also has an S_3-symmetry. In this paper, we prove an S_3-symmetry of the Jacobi identity for intertwining operator algebras introduced by Huang [H2] and studied by the author [C]. See Theorem 3.1 for the statement of this S_3-symmetry. The S_3-symmetry in this general case is much more complicated but is also much more interesting and much deeper. Note that the Jacobi identity for general intertwining operator algebras in [H2] and [C] involves the braiding and fusing isomorphisms satisfying the genus-zero Moore-Seiberg equations. The proof of the S_3-symmetry in the present paper uses not only the properties of the intertwining operators (for example, the skew-symmetry) but also the properties of braiding and fusing isomorphisms and the genus-zero Moore-Seiberg equations. In particular, our proof depends heavily on the theory of multivalued analytic functions of several variables, especially the theory of analytic extensions.

This paper is organized as follows. In section 2, we review some preliminaries concerning the theory of intertwining operator algebras which we need to formulate and prove the main result of this paper. In section 3, we prove an S_3-symmetry of the Jacobi identity for intertwining operator algebras.

Acknowledgments The author is very grateful to Professor Yi-Zhi Huang for his support, encouragement, many discussions on the paper and help with the exposition of the paper. The author is supported by NSFC grant 11401559.

2 Preliminaries

In this section, we first recall some notations and facts in formal calculus and complex analysis (see [FLM, FHL, H2] for more details), then we review some definitions and properties in the theory of intertwining operator algebras in [H2, C]. These are necessary preliminaries for formulating and proving the main result of this paper.

In this paper, as in [FHL, H2, C], x, $x_0, \dots$ are independent commuting formal variables. And for a vector space W and a formal variable x, as in [FHL, H2, C], we shall use $W[x]$, $W[x, x^{-1}]$, $W[[x]]$, $W[[x, x^{-1}]]$, $W((x))$ and $W\{x\}$ to denote the spaces of all polynomials in x, all Laurent polynomials in x, all formal power series in x, all formal Laurent series in x, all formal Laurent series in x with finitely many negative powers and all formal series with arbitrary powers of x in $\mathbb{C}$, respectively. For series with more than one formal variables, we shall use similar notations. We shall use $\mathrm{Res}_x f(x)$ to denote the coefficient of x^{-1} in $f(x)$ for any $f(x) \in W\{x\}$. As in [FHL, H2, C], $z, z_0, \dots,$ are complex numbers, *not* formal variables.

Let

$$\delta(x) = \sum_{n\in\mathbb{Z}} x^n. \tag{2.1}$$

It has the following important property: For any $f(x) \in \mathbb{C}[x, x^{-1}]$,

$$f(x)\delta(x) = f(1)\delta(x). \tag{2.2}$$

Following [FHL, H2, C], we use the convention that negative powers of a binomial are to be expanded in nonnegative powers of the second summand so that, for example,

$$x_0^{-1}\delta\left(\frac{x_1 - x_2}{x_0}\right) = \sum_{n\in\mathbb{Z}} \frac{(x_1 - x_2)^n}{x_0^{n+1}} = \sum_{m\in\mathbb{N},\, n\in\mathbb{Z}} (-1)^m \binom{n}{m} x_0^{-n-1} x_1^{n-m} x_2^m. \tag{2.3}$$

The following identities are often very useful:

$$x_1^{-1}\delta\left(\frac{x_2+x_0}{x_1}\right)=x_2^{-1}\delta\left(\frac{x_1-x_0}{x_2}\right), \tag{2.4}$$

$$x_0^{-1}\delta\left(\frac{x_1-x_2}{x_0}\right)-x_0^{-1}\delta\left(\frac{x_2-x_1}{-x_0}\right)=x_2^{-1}\delta\left(\frac{x_1-x_0}{x_2}\right). \tag{2.5}$$

As in [FHL, H2, C], $\mathbb{C}[x_1, x_2]_S$ is the ring of rational functions obtained by inverting the products of (zero or more) elements of the set S of nonzero homogenous linear polynomials in x_1 and x_2. Also, ι_{12} is the operation of expanding an element of $\mathbb{C}[x_1, x_2]_S$, that is, a polynomial in x_1 and x_2 divided by a product of homogenous linear polynomials in x_1 and x_2, as a formal series containing at most finitely many negative powers of x_2 (using binomial expansions for negative powers of linear polynomials involving both x_1 and x_2); similarly for ι_{21}, and so on. We need the following fact from [FHL].

Proposition 2.1. *Consider a rational function of the form*

$$f(x_0,x_1,x_2)=\frac{g(x_0,x_1,x_2)}{x_0^r x_1^s x_2^t}, \tag{2.6}$$

where g is a polynomial and $r, s, t \in \mathbb{Z}$. Then

$$x_1^{-1}\delta\left(\frac{x_2+x_0}{x_1}\right)\iota_{20}(f|_{x_1=x_0+x_2})=x_2^{-1}\delta\left(\frac{x_1-x_0}{x_2}\right)\iota_{10}(f|_{x_2=x_1-x_0}) \tag{2.7}$$

and

$$\begin{aligned}x_0^{-1}\delta\left(\frac{x_1-x_2}{x_0}\right)\iota_{12}(f|_{x_0=x_1-x_2})-x_0^{-1}\delta\left(\frac{x_2-x_1}{-x_0}\right)\iota_{21}(f|_{x_0=x_1-x_2})\\ =x_2^{-1}\delta\left(\frac{x_1-x_0}{x_2}\right)\iota_{10}(f|_{x_2=x_1-x_0}).\end{aligned} \tag{2.8}$$

As in [FHL, H2, C], the graded dual of a $\mathbb{Z}$-graded, or more generally, $\mathbb{C}$-graded, vector space $W = \coprod_n W_{(n)}$ is denoted by

$$W' = \coprod_n W_{(n)}^*. \tag{2.9}$$

For any $z \in \mathbb{C}$, we use $\log z$ to denote the value $\log|z| + i \arg z$ with $0 \le \arg z < 2\pi$ of logarithm of z. For two multivalued functions f_1 and f_2 on a region, f_1 and f_2 are equal if on any simply connected open subset of the region, any single-valued branch of f_1 is equal to a single-valued branch of f_2, and vice versa.

Now we recall some basic notions and results in the theory of intertwining operator algebras. For the details of the definitions and properties of vertex operator algebras, their modules and intertwining operators, the reader is referred to [FHL, FLM, H2]. And for more details of the properties of intertwining operator algebras, the reader is referred to [H2, C].

Let $(V, Y, \mathbf{1}, \omega)$ be a vertex operator algebra, and let W_1, W_2, W_3 be modules of V. The space of all intertwining operators of type $\binom{W_3}{W_1\ W_2}$ is denoted by $\bar{\mathcal{V}}^{W_3}_{W_1W_2}$ instead of $\mathcal{V}^{W_3}_{W_1W_2}$, for as in [C], the latter shall be used to denote a subspace of $\bar{\mathcal{V}}^{W_3}_{W_1W_2}$ in the definition of intertwining operator algebra. The dimension of this vector space is denoted by $\bar{\mathcal{N}}^{W_3}_{W_1W_2}$. It is the so-called fusion rule of the same type. Let $\mathcal{Y}$ be an intertwining operator of type $\binom{W_3}{W_1W_2}$. Given any $r \in \mathbb{Z}$, as in [HL, H2, C], we define

$$\Omega_r(\mathcal{Y}) : W_2 \otimes W_1 \to W_3\{x\} \tag{2.10}$$

by

$$\Omega_r(\mathcal{Y})(w_{(2)}, x)w_{(1)} = e^{xL(-1)}\mathcal{Y}(w_{(1)}, e^{(2r+1)\pi i}x)w_{(2)} \tag{2.11}$$

for $w_{(1)} \in W_1$, $w_{(2)} \in W_2$. We have the following result proved in [HL]:

Proposition 2.2. *For any* $\mathcal{Y} \in \bar{\mathcal{V}}^{W_3}_{W_1W_2}$, $r \in \mathbb{Z}$, *we have* $\Omega_r(\mathcal{Y}) \in \bar{\mathcal{V}}^{W_3}_{W_2W_1}$. *Moreover,*

$$\Omega_{-r-1}(\Omega_r(\mathcal{Y})) = \Omega_r(\Omega_{-r-1}(\mathcal{Y})) = \mathcal{Y}. \tag{2.12}$$

In particular, the correspondence $\mathcal{Y} \mapsto \Omega_r(\mathcal{Y})$ *defines a linear isomorphism from* $\bar{\mathcal{V}}^{W_3}_{W_1W_2}$ *to* $\bar{\mathcal{V}}^{W_3}_{W_2W_1}$, *and we have*

$$\bar{\mathcal{N}}^{W_3}_{W_1W_2} = \bar{\mathcal{N}}^{W_3}_{W_2W_1}. \tag{2.13}$$

Now we recall the first definition of intertwining operator algebras in [H2]:

Definition 2.3 (Intertwining operator algebra). An *intertwining operator algebra of central charge* $c \in \mathbb{C}$ consists of the following data:

1. A vector space
$$W = \coprod_{a\in\mathcal{A}} W^a \tag{2.14}$$
graded by a finite set $\mathcal{A}$ containing a special element e (graded by *color*).

2. A vertex operator algebra structure of central charge c on W^e, and a W^e-module structure on W^a for each $a \in \mathcal{A}$.

3. A subspace $\mathcal{V}^{a_3}_{a_1a_2}$ of the space of all intertwining operators of type $\binom{W^{a_3}}{W^{a_1}W^{a_2}}$ for each triple $a_1, a_2, a_3 \in \mathcal{A}$, with its dimension denoted by $\mathcal{N}^{a_3}_{a_1a_2}$.

These data satisfy the following axioms for any $a_1, a_2, a_3, a_4, a_5, a_6 \in \mathcal{A}$, $w_{(a_i)} \in W^{a_i}$, $i = 1, 2, 3$, and $w'_{(a_4)} \in (W^{a_4})'$:

1. The W^e-module structure on W^e is the adjoint module structure. For any $a \in \mathcal{A}$, the space $\mathcal{V}_{ea}^{a}$ is the one-dimensional vector space spanned by the vertex operator for the W^e-module W^a. For any $a_1, a_2 \in \mathcal{A}$ such that $a_1 \neq a_2$, $\mathcal{V}_{ea_1}^{a_2} = 0$.

2. *Weight condition*: For any $a \in \mathcal{A}$ and the corresponding module $W^a = \coprod_{n\in\mathbb{C}} W^a_{(n)}$ graded by the action of $L(0)$, there exists $h_a \in \mathbb{R}$ such that $W^a_{(n)} = 0$ for $n \notin h_a + \mathbb{Z}$.

3. *Convergence properties*: For any $m \in \mathbb{Z}_+$, $a_i, b_j \in \mathcal{A}$, $w_{(a_i)} \in W^{a_i}$, $\mathcal{Y}_i \in \mathcal{V}^{b_i}_{a_i\, b_{i+1}}$, $i = 1, \dots, m$, $j = 1, \dots, m+1$, $w'_{(b_1)} \in (W^{b_1})'$ and $w_{(b_{m+1})} \in W^{b_{m+1}}$, the series

$$\langle w'_{(b_1)}, \mathcal{Y}_1(w_{(a_1)}, x_1) \cdots \mathcal{Y}_m(w_{(a_m)}, x_m) w_{(b_{m+1})}\rangle_{W^{b_1}}\big|_{x_i^n = e^{n \log z_i},\ i=1,\dots,m,\ n\in\mathbb{R}} \tag{2.15}$$

is absolutely convergent when $|z_1| > \cdots > |z_m| > 0$ and its sum can be analytically extended to a multivalued analytic function on the region given by $z_i \neq 0$, $i = 1, \dots, m$, $z_i \neq z_j$, $i \neq j$, such that for any set of possible singular points with either $z_i = 0$, $z_i = \infty$ or $z_i = z_j$ for $i \neq j$, this multivalued analytic function can be expanded near the singularity as a series having the same form as the expansion near the singular points of a solution of a system of differential equations with regular singular points. For any $\mathcal{Y}_1 \in \mathcal{V}^{a_5}_{a_1 a_2}$ and $\mathcal{Y}_2 \in \mathcal{V}^{a_4}_{a_5 a_3}$, the series

$$\langle w'_{(a_4)}, \mathcal{Y}_2(\mathcal{Y}_1(w_{(a_1)}, x_0) w_{(a_2)}, x_2) w_{(a_3)}\rangle_{W^{a_4}}\big|_{x_0^n = e^{n\log(z_1 - z_2)},\ x_2^n = e^{n \log z_2},\ n\in\mathbb{R}} \tag{2.16}$$

is absolutely convergent when $|z_2| > |z_1 - z_2| > 0$.

4. *Associativity*: For any $\mathcal{Y}_1 \in \mathcal{V}^{a_4}_{a_1 a_5}$ and $\mathcal{Y}_2 \in \mathcal{V}^{a_5}_{a_2 a_3}$, there exist $\mathcal{Y}^a_{3,i} \in \mathcal{V}^a_{a_1 a_2}$ and $\mathcal{Y}^a_{4,i} \in \mathcal{V}^{a_4}_{a a_3}$ for $i = 1, \dots, \mathcal{N}^a_{a_1 a_2}\mathcal{N}^{a_4}_{a a_3}$ and $a \in \mathcal{A}$, such that the (multivalued) analytic function

$$\langle w'_{(a_4)}, \mathcal{Y}_1(w_{(a_1)}, x_1)\mathcal{Y}_2(w_{(a_2)}, x_2) w_{(a_3)}\rangle_{W^{a_4}}\big|_{x_1 = z_1, x_2 = z_2} \tag{2.17}$$

defined on the region $|z_1| > |z_2| > 0$ and the (multivalued) analytic function

$$\sum_{a\in\mathcal{A}} \sum_{i=1}^{\mathcal{N}^a_{a_1a_2}\mathcal{N}^{a_4}_{aa_3}} \langle w'_{(a_4)}, \mathcal{Y}^a_{4,i}(\mathcal{Y}^a_{3,i}(w_{(a_1)}, x_0) w_{(a_2)}, x_2) w_{(a_3)}\rangle_{W^{a_4}}\Big|_{x_0 = z_1 - z_2, x_2 = z_2} \tag{2.18}$$

defined on the region $|z_2| > |z_1 - z_2| > 0$ are equal on the intersection $|z_1| > |z_2| > |z_1 - z_2| > 0$.

5. *Skew-symmetry*: The restriction of Ω_{-1} to $\mathcal{V}^{a_3}_{a_1 a_2}$ is an isomorphism from $\mathcal{V}^{a_3}_{a_1 a_2}$ to $\mathcal{V}^{a_3}_{a_2 a_1}$.

Remark 2.4. *The skew-symmetry isomorphisms* $\Omega_{-1}(a_1, a_2; a_3)$ *for all* $a_1, a_2, a_3 \in \mathcal{A}$ *give an isomorphism*

$$\Omega_{-1} : \coprod_{a_1,a_2,a_3\in\mathcal{A}} \mathcal{V}^{a_3}_{a_1 a_2} \to \coprod_{a_1,a_2,a_3\in\mathcal{A}} \mathcal{V}^{a_3}_{a_1 a_2}, \tag{2.19}$$

which, as in [H2, C], is still called the skew-symmetry isomorphism. In this paper, as in [H2, C], we shall omit subscript -1 *in* Ω_{-1} *for simplicity and denote it by* Ω.

We denote the intertwining operator algebra just defined by $(W, \mathcal{A}, \{\mathcal{V}_{a_1a_2}^{a_3}\}, \mathbf{1}, \omega)$ or simply by W.

Next, as in [H2, C], we give the two linear maps corresponding to the multiplication and iterates of intertwining operators, respectively. Let

$$\begin{aligned} \mathbf{P} : \coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5} &\to (\mathrm{Hom}(W\otimes W\otimes W, W))\{x_1, x_2\} \\ \mathcal{Z} &\mapsto \mathbf{P}(\mathcal{Z}) \end{aligned} \tag{2.20}$$

be the linear map defined using products of intertwining operators as follows: For

$$\mathcal{Z} \in \coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5}, \tag{2.21}$$

the element $\mathbf{P}(\mathcal{Z})$ to be defined is a linear map from $W\otimes W\otimes W$ to $W\{x_1, x_2\}$. We denote the image of $w_1\otimes w_2\otimes w_3$ under this map by $(\mathbf{P}(\mathcal{Z}))(w_1, w_2, w_3; x_1, x_2)$ for any $w_1, w_2, w_3 \in W$. Then we define $\mathbf{P}$ by linearity and by

$$\begin{aligned} &(\mathbf{P}(\mathcal{Y}_1\otimes\mathcal{Y}_2))(w_{(a_6)}, w_{(a_7)}, w_{(a_8)}; x_1, x_2) \\ &\quad = \begin{cases} \mathcal{Y}_1(w_{(a_6)}, x_1)\mathcal{Y}_2(w_{(a_7)}, x_2)w_{(a_8)}, & a_6 = a_1, a_7 = a_2, a_8 = a_3, \\ 0, & \text{otherwise} \end{cases} \end{aligned} \tag{2.22}$$

for $a_1, \dots, a_8 \in \mathcal{A}$, $\mathcal{Y}_1 \in \mathcal{V}_{a_1a_5}^{a_4}$, $\mathcal{Y}_2 \in \mathcal{V}_{a_2a_3}^{a_5}$, and $w_{(a_6)} \in W^{a_6}$, $w_{(a_7)} \in W^{a_7}$, $w_{(a_8)} \in W^{a_8}$. So we have an isomorphism

$$\tilde{\mathbf{P}} : \frac{\coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5}}{\mathrm{Ker}\ \mathbf{P}} \longrightarrow \mathbf{P}\left(\coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5}\right) \tag{2.23}$$

which makes the following diagram commute:

$$\begin{tikzcd} \coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5} \arrow[r, "\mathbf{P}"] \arrow[d, "\pi_P"'] & \mathbf{P}\left(\coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5}\right), \\ \frac{\coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5}}{\mathrm{Ker}\ \mathbf{P}} \arrow[ur, "\tilde{\mathbf{P}}"'] & \end{tikzcd} \tag{2.24}$$

where π_P is the corresponding canonical projective map. As in [C], we also denote $\pi_P(\mathcal{Z})$ by $[\mathcal{Z}]_P$ or $\mathcal{Z} + \mathrm{Ker}\ \mathbf{P}$ for $\mathcal{Z} \in \coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5}$ when there is no ambiguity. The second linear map is

$$\begin{aligned} \mathbf{I} : \coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_2}^{a_5} \otimes \mathcal{V}_{a_5a_3}^{a_4} &\to (\mathrm{Hom}(W\otimes W\otimes W, W))\{x_0, x_2\} \\ \mathcal{Z} &\mapsto \mathbf{I}(\mathcal{Z}) \end{aligned} \tag{2.25}$$

defined using iterates of intertwining operators as follows: For

$$\mathcal{Z} \in \coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_2}^{a_5} \otimes \mathcal{V}_{a_5a_3}^{a_4}, \tag{2.26}$$

the element $\mathbf{I}(\mathcal{Z})$ to be defined is a linear map from $W\otimes W\otimes W$ to $W\{x_0, x_2\}$. We denote the image of $w_1 \otimes w_2 \otimes w_3$ under this map by $(\mathbf{I}(\mathcal{Z}))(w_1, w_2, w_3; x_0, x_2)$ for any $w_1, w_2, w_3 \in W$. Then we define $\mathbf{I}$ by linearity and by

$$\begin{aligned}
&(\mathbf{I}(\mathcal{Y}_1 \otimes \mathcal{Y}_2))(w_{(a_6)}, w_{(a_7)}, w_{(a_8)}; x_0, x_2)\\
&\quad= \begin{cases} \mathcal{Y}_2(\mathcal{Y}_1(w_{(a_6)}, x_0)w_{(a_7)}, x_2)w_{(a_8)}, & a_6 = a_1, a_7 = a_2, a_8 = a_3,\\ 0, & \text{otherwise} \end{cases}
\end{aligned} \tag{2.27}$$

for $a_1, \dots, a_8 \in \mathcal{A}$, $\mathcal{Y}_1 \in \mathcal{V}_{a_1a_2}^{a_5}$, $\mathcal{Y}_2 \in \mathcal{V}_{a_5a_3}^{a_4}$, and $w_{(a_6)} \in W^{a_6}$, $w_{(a_7)} \in W^{a_7}$, $w_{(a_8)} \in W^{a_8}$. Therefore we have an isomorphism

$$\tilde{\mathbf{I}} : \quad \frac{\displaystyle\coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_2}^{a_5} \otimes \mathcal{V}_{a_5a_3}^{a_4}}{\operatorname{Ker} \mathbf{I}} \longrightarrow \mathbf{I}\left(\coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_2}^{a_5} \otimes \mathcal{V}_{a_5a_3}^{a_4}\right) \tag{2.28}$$

which makes the following diagram commute:

$$\begin{tikzcd}
\displaystyle\coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_2}^{a_5} \otimes \mathcal{V}_{a_5a_3}^{a_4} \arrow[r, "\mathbf{I}"] \arrow[d, "\pi_I"'] & \mathbf{I}\left(\displaystyle\coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_2}^{a_5} \otimes \mathcal{V}_{a_5a_3}^{a_4}\right), \\
\dfrac{\displaystyle\coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_2}^{a_5} \otimes \mathcal{V}_{a_5a_3}^{a_4}}{\operatorname{Ker} \mathbf{I}} \arrow[ur, "\tilde{\mathbf{I}}"] &
\end{tikzcd} \tag{2.29}$$

where π_I is the corresponding canonical projective map. As in [C], we also denote $\pi_I(\mathcal{Z})$ by $[\mathcal{Z}]_I$ or $\mathcal{Z} + \operatorname{Ker} \mathbf{I}$ for $\mathcal{Z} \in \coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_2}^{a_5} \otimes \mathcal{V}_{a_5a_3}^{a_4}$ when there is no ambiguity.

The two linear maps $\mathbf{P}$ and $\mathbf{I}$ are called the *multiplication of intertwining operators* and the *iterates of intertwining operators*, respectively.

Moreover, in [H2, C], Huang and the author obtained isomorphisms from the associativity of intertwining operator algebras and from the skew-symmetry isomorphism Ω. The *fusing isomorphism* which we obtained from the associativity of intertwining operator algebras is a map

$$\mathcal{F} : \quad \frac{\displaystyle\coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5}}{\operatorname{Ker} \mathbf{P}} \longrightarrow \frac{\displaystyle\coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_2}^{a_5} \otimes \mathcal{V}_{a_5a_3}^{a_4}}{\operatorname{Ker} \mathbf{I}} \tag{2.30}$$

determined by linearity and by

$$\mathcal{F}(\mathcal{Y}_1 \otimes \mathcal{Y}_2 + \text{Ker } \mathbf{P}) = \sum_{a \in \mathcal{A}} \sum_{i=1}^{\mathcal{N}_{a_1a_2}^{a}\mathcal{N}_{aa_3}^{a_4}} \mathcal{Y}_{3,i}^{a} \otimes \mathcal{Y}_{4,i}^{a} + \text{Ker } \mathbf{I} \tag{2.31}$$

for $a_1, \cdots, a_5 \in \mathcal{A}$, $\mathcal{Y}_1 \in \mathcal{V}_{a_1a_5}^{a_4}$ and $\mathcal{Y}_2 \in \mathcal{V}_{a_2a_3}^{a_5}$, where

$$\{\mathcal{Y}_{3,i}^{a} \in \mathcal{V}_{a_1a_2}^{a}, \mathcal{Y}_{4,i}^{a} \in \mathcal{V}_{aa_3}^{a_4} \mid i = 1, \cdots, \mathcal{N}_{a_1a_2}^{a}\mathcal{N}_{aa_3}^{a_4}, a \in \mathcal{A}\} \tag{2.32}$$

is a set of intertwining operators satisfying that for any $w_1, w_2, w_3 \in W$ and $w' \in W'$,

$$\sum_{a \in \mathcal{A}} \sum_{i=1}^{\mathcal{N}_{a_1a_2}^{a}\mathcal{N}_{aa_3}^{a_4}} \langle w', (\mathbf{I}(\mathcal{Y}_{3,i}^{a} \otimes \mathcal{Y}_{4,i}^{a}))(w_1, w_2, w_3; x_0, x_2)\rangle_W \Big|_{x_0^n = e^{n \log(z_1 - z_2)}, x_2^n = e^{n \log z_2}} \tag{2.33}$$

is equal to

$$\langle w', (\mathbf{P}(\mathcal{Y}_1 \otimes \mathcal{Y}_2))(w_1, w_2, w_3; x_1, x_2)\rangle_W \big|_{x_1^n = e^{n \log z_1}, x_2^n = e^{n \log z_2}} \tag{2.34}$$

on the region

$$S_1 = \{(z_1, z_2) \in \mathbb{C}^2 \mid \text{Re}z_1 > \text{Re}z_2 > \text{Re}(z_1 - z_2) > 0,\ \text{Im}z_1 > \text{Im}z_2 > \text{Im}(z_1 - z_2) > 0\}.$$

It was also proved that

$$\begin{aligned} \mathcal{F}(a_1, a_2, a_3, a_4): \quad \pi_P\left(\coprod_{a_5 \in \mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5}\right) &\longrightarrow \pi_I\left(\coprod_{a_5 \in \mathcal{A}} \mathcal{V}_{a_1a_2}^{a_5} \otimes \mathcal{V}_{a_5a_3}^{a_4}\right) \\ \mathcal{Y}_1 \otimes \mathcal{Y}_2 + \text{Ker } \mathbf{P} &\longmapsto \mathcal{F}(\mathcal{Y}_1 \otimes \mathcal{Y}_2 + \text{Ker } \mathbf{P}) \end{aligned} \tag{2.35}$$

is an isomorphism for any $a_1, \cdots, a_4 \in \mathcal{A}$, where $\mathcal{Y}_1 \in \mathcal{V}_{a_1a_5}^{a_4}$, $\mathcal{Y}_2 \in \mathcal{V}_{a_2a_3}^{a_5}$. These isomorphisms are also called *fusing isomorphisms*. The isomorphisms we obtained from Ω and its inverse are linear isomorphic maps:

$$\tilde{\Omega}^{(1)}, (\widetilde{\Omega^{-1}})^{(1)}: \frac{\coprod_{a_1,a_2,a_3,a_4,a_5 \in \mathcal{A}} \mathcal{V}_{a_1a_2}^{a_5} \otimes \mathcal{V}_{a_5a_3}^{a_4}}{\text{Ker } \mathbf{I}} \longrightarrow \frac{\coprod_{a_1,a_2,a_3,a_4,a_5 \in \mathcal{A}} \mathcal{V}_{a_2a_1}^{a_5} \otimes \mathcal{V}_{a_5a_3}^{a_4}}{\text{Ker } \mathbf{I}} \tag{2.36}$$

defined by linearity and by

$$\tilde{\Omega}^{(1)}(\mathcal{Y}_1 \otimes \mathcal{Y}_2 + \text{Ker } \mathbf{I}) = \Omega(\mathcal{Y}_1) \otimes \mathcal{Y}_2 + \text{Ker } \mathbf{I}, \tag{2.37}$$

$$(\widetilde{\Omega^{-1}})^{(1)}(\mathcal{Y}_1 \otimes \mathcal{Y}_2 + \text{Ker } \mathbf{I}) = \Omega^{-1}(\mathcal{Y}_1) \otimes \mathcal{Y}_2 + \text{Ker } \mathbf{I} \tag{2.38}$$

for $a_1, \ldots, a_5 \in \mathcal{A}$, $\mathcal{Y}_1 \in \mathcal{V}_{a_1a_2}^{a_5}$, $\mathcal{Y}_2 \in \mathcal{V}_{a_5a_3}^{a_4}$;

$$\tilde{\Omega}^{(2)}, (\widetilde{\Omega^{-1}})^{(2)}: \frac{\coprod_{a_1,a_2,a_3,a_4,a_5 \in \mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5}}{\text{Ker } \mathbf{P}} \longrightarrow \frac{\coprod_{a_1,a_2,a_3,a_4,a_5 \in \mathcal{A}} \mathcal{V}_{a_2a_3}^{a_5} \otimes \mathcal{V}_{a_5a_1}^{a_4}}{\text{Ker } \mathbf{I}} \tag{2.39}$$

defined by linearity and by

$$\tilde{\Omega}^{(2)}(\mathcal{Y}_1 \otimes \mathcal{Y}_2 + \mathrm{Ker}\,\mathbf{P}) = \mathcal{Y}_2 \otimes \Omega(\mathcal{Y}_1) + \mathrm{Ker}\,\mathbf{I}, \tag{2.40}$$

$$(\widetilde{\Omega^{-1}})^{(2)}(\mathcal{Y}_1 \otimes \mathcal{Y}_2 + \mathrm{Ker}\,\mathbf{P}) = \mathcal{Y}_2 \otimes \Omega^{-1}(\mathcal{Y}_1) + \mathrm{Ker}\,\mathbf{I} \tag{2.41}$$

for $a_1, \dots, a_5 \in \mathcal{A}$, $\mathcal{Y}_1 \in \mathcal{V}_{a_1a_5}^{a_4}$, $\mathcal{Y}_2 \in \mathcal{V}_{a_2a_3}^{a_5}$;

$$\tilde{\Omega}^{(3)}, (\widetilde{\Omega^{-1}})^{(3)} : \frac{\coprod_{a_1,a_2,a_3,a_4,a_5 \in \mathcal{A}} \mathcal{V}_{a_1a_2}^{a_5} \otimes \mathcal{V}_{a_5a_3}^{a_4}}{\mathrm{Ker}\,\mathbf{I}} \longrightarrow \frac{\coprod_{a_1,a_2,a_3,a_4,a_5 \in \mathcal{A}} \mathcal{V}_{a_3a_5}^{a_4} \otimes \mathcal{V}_{a_1a_2}^{a_5}}{\mathrm{Ker}\,\mathbf{P}} \tag{2.42}$$

defined by linearity and by

$$\tilde{\Omega}^{(3)}(\mathcal{Y}_1 \otimes \mathcal{Y}_2 + \mathrm{Ker}\,\mathbf{I}) = \Omega(\mathcal{Y}_2) \otimes \mathcal{Y}_1 + \mathrm{Ker}\,\mathbf{P}, \tag{2.43}$$

$$(\widetilde{\Omega^{-1}})^{(3)}(\mathcal{Y}_1 \otimes \mathcal{Y}_2 + \mathrm{Ker}\,\mathbf{I}) = \Omega^{-1}(\mathcal{Y}_2) \otimes \mathcal{Y}_1 + \mathrm{Ker}\,\mathbf{P} \tag{2.44}$$

for $a_1, \dots, a_5 \in \mathcal{A}$, $\mathcal{Y}_1 \in \mathcal{V}_{a_1a_2}^{a_5}$, $\mathcal{Y}_2 \in \mathcal{V}_{a_5a_3}^{a_4}$;

$$\tilde{\Omega}^{(4)}, (\widetilde{\Omega^{-1}})^{(4)} : \frac{\coprod_{a_1,a_2,a_3,a_4,a_5 \in \mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5}}{\mathrm{Ker}\,\mathbf{P}} \longrightarrow \frac{\coprod_{a_1,a_2,a_3,a_4,a_5 \in \mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_3a_2}^{a_5}}{\mathrm{Ker}\,\mathbf{P}} \tag{2.45}$$

defined by linearity and by

$$\tilde{\Omega}^{(4)}(\mathcal{Y}_1 \otimes \mathcal{Y}_2 + \mathrm{Ker}\,\mathbf{P}) = \mathcal{Y}_1 \otimes \Omega(\mathcal{Y}_2) + \mathrm{Ker}\,\mathbf{P}, \tag{2.46}$$

$$(\widetilde{\Omega^{-1}})^{(4)}(\mathcal{Y}_1 \otimes \mathcal{Y}_2 + \mathrm{Ker}\,\mathbf{P}) = \mathcal{Y}_1 \otimes \Omega^{-1}(\mathcal{Y}_2) + \mathrm{Ker}\,\mathbf{P} \tag{2.47}$$

for $a_1, \dots, a_5 \in \mathcal{A}$, $\mathcal{Y}_1 \in \mathcal{V}_{a_1a_5}^{a_4}$, $\mathcal{Y}_2 \in \mathcal{V}_{a_2a_3}^{a_5}$. And these isomorphisms have relations:

$$(\tilde{\Omega}^{(2)})^{-1} = (\widetilde{\Omega^{-1}})^{(3)}, \quad ((\widetilde{\Omega^{-1}})^{(2)})^{-1} = \tilde{\Omega}^{(3)}, \tag{2.48}$$

$$(\tilde{\Omega}^{(1)})^{-1} = (\widetilde{\Omega^{-1}})^{(1)}, \quad (\tilde{\Omega}^{(4)})^{-1} = (\widetilde{\Omega^{-1}})^{(4)}. \tag{2.49}$$

The above isomorphisms are not independent, we proved the following relations in [C]:

Theorem 2.5. *The above isomorphisms satisfy the following genus-zero Moore-Seiberg equations:*

$$\begin{aligned} \mathcal{F} \circ \tilde{\Omega}^{(3)} \circ \mathcal{F} &= \tilde{\Omega}^{(1)} \circ \mathcal{F} \circ \tilde{\Omega}^{(4)}, \qquad &(2.50)\\ \mathcal{F} \circ (\widetilde{\Omega^{-1}})^{(3)} \circ \mathcal{F} &= (\widetilde{\Omega^{-1}})^{(1)} \circ \mathcal{F} \circ (\widetilde{\Omega^{-1}})^{(4)}. \qquad &(2.51) \end{aligned}$$

Using the fusing isomorphism and the isomorphism $\tilde{\Omega}^{(1)}$, we deduced a *braiding isomorphism*

$$\mathcal{B} = \mathcal{F}^{-1} \circ \tilde{\Omega}^{(1)} \circ \mathcal{F} : \frac{\coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5}}{\text{Ker } \mathbf{P}} \longrightarrow \frac{\coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_2a_5}^{a_4} \otimes \mathcal{V}_{a_1a_3}^{a_5}}{\text{Ker } \mathbf{P}}. \tag{2.52}$$

Moreover, we get an isomorphism

$$\begin{aligned} \mathcal{B}(a_1,a_2,a_3,a_4) : \quad \pi_P\left(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5}\right) &\longrightarrow \pi_P\left(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_2a_5}^{a_4} \otimes \mathcal{V}_{a_1a_3}^{a_5}\right) \\ \mathcal{Y}_1 \otimes \mathcal{Y}_2 + \text{Ker } \mathbf{P} &\longmapsto \mathcal{B}(\mathcal{Y}_1 \otimes \mathcal{Y}_2 + \text{Ker } \mathbf{P}) \end{aligned} \tag{2.53}$$

for any $a_1, \cdots, a_4 \in \mathcal{A}$, where $\mathcal{Y}_1 \in \mathcal{V}_{a_1a_5}^{a_4}$, $\mathcal{Y}_2 \in \mathcal{V}_{a_2a_3}^{a_5}$. We also call these isomorphisms *braiding isomorphisms.*

Before formulating the Jacobi identity for intertwining operator algebras, we need to recall the specifics of one more property, which is about certain special multivalued analytic functions, and were discussed in [H2, C].

First we consider some simply connected regions in $\mathbb{C}^2$. Cutting the regions $|z_1| > |z_2| > 0$ and $|z_2| > |z_1| > 0$ along the intersections of these regions with

$$\{(z_1, z_2) \in \mathbb{C}^2 \mid z_1 \in [0, +\infty)\} \cup \{(z_1, z_2) \in \mathbb{C}^2 \mid z_2 \in [0, +\infty)\},$$

we obtain two simply connected regions, which, as in [H2, C], are denoted by R_1 and R_2, respectively. Also, let R_3 be the simply connected region obtained by cutting the region $|z_2| > |z_1 - z_2| > 0$ along the intersection of this region with

$$\{(z_1, z_2) \in \mathbb{C}^2 \mid z_2 \in [0, +\infty)\} \cup \{(z_1, z_2) \in \mathbb{C}^2 \mid z_1 - z_2 \in [0, +\infty)\},$$

and let R_4 be the simply connected region obtained by cutting the region $|z_1| > |z_1 - z_2| > 0$ along the intersection of this region with

$$\{(z_1, z_2) \in \mathbb{C}^2 \mid z_1 \in [0, +\infty)\} \cup \{(z_1, z_2) \in \mathbb{C}^2 \mid z_2 - z_1 \in [0, +\infty)\}.$$

Then we consider some special multivalued analytic functions on

$$M^2 = \{(z_1, z_2) \in \mathbb{C}^2 \mid z_1, z_2 \neq 0, z_1 \neq z_2\}. \tag{2.54}$$

For $a_1, a_2, a_3, a_4 \in \mathcal{A}$, as in [H2, C], we let $\mathbb{G}^{a_1,a_2,a_3,a_4}$ be the set of multivalued analytic functions on M^2 with a choice of a single-valued branch on the region R_1 satisfying the following property: Any branch of $f(z_1, z_2) \in \mathbb{G}^{a_1,a_2,a_3,a_4}$ on the regions $|z_1| > |z_2| > 0$, $|z_2| > |z_1| > 0$ and $|z_2| > |z_1 - z_2| > 0$, respectively, can be expanded as

$$\sum_{a\in\mathcal{A}} z_1^{h_{a_4}-h_{a_1}-h_a} z_2^{h_a-h_{a_2}-h_{a_3}} F_a(z_1, z_2), \tag{2.55}$$

$$\sum_{a\in\mathcal{A}} z_2^{h_{a_4}-h_{a_2}-h_a} z_1^{h_a-h_{a_1}-h_{a_3}} G_a(z_1, z_2) \tag{2.56}$$

and

$$\sum_{a\in\mathcal{A}} z_2^{h_{a_4}-h_a-h_{a_3}} (z_1 - z_2)^{h_a-h_{a_1}-h_{a_2}} H_a(z_1, z_2), \tag{2.57}$$

respectively, where for $a \in \mathcal{A}$,

$$F_a(z_1, z_2) \in \mathbb{C}[[z_2/z_1]][z_1, z_1^{-1}, z_2, z_2^{-1}], \tag{2.58}$$

$$G_a(z_1, z_2) \in \mathbb{C}[[z_1/z_2]][z_1, z_1^{-1}, z_2, z_2^{-1}] \tag{2.59}$$

and

$$H_a(z_1, z_2) \in \mathbb{C}[[(z_1 - z_2)/z_2]][z_2, z_2^{-1}, z_1 - z_2, (z_1 - z_2)^{-1}]. \tag{2.60}$$

The chosen single-valued branch on R_1 of an element of $\mathbb{G}^{a_1,a_2,a_3,a_4}$ is called the *preferred branch on* R_1. As in [C], we use the nonempty simply connected regions

$$S_1 = \{(z_1, z_2) \in \mathbb{C}^2 \mid \mathrm{Re} z_1 > \mathrm{Re} z_2 > \mathrm{Re}(z_1 - z_2) > 0,\ \mathrm{Im} z_1 > \mathrm{Im} z_2 > \mathrm{Im}(z_1 - z_2) > 0\}$$

and

$$S_2 = \{(z_1, z_2) \in \mathbb{C}^2 \mid \mathrm{Re} z_2 > \mathrm{Re} z_1 > \mathrm{Re}(z_2 - z_1) > 0,\ \mathrm{Im} z_2 > \mathrm{Im} z_1 > \mathrm{Im}(z_2 - z_1) > 0\}$$

to determine other special branches of an element of $\mathbb{G}^{a_1,a_2,a_3,a_4}$ on R_2, R_3 and R_4 related to the preferred branch on R_1. Firstly, the restriction of the preferred branch on R_1 of an element of $\mathbb{G}^{a_1,a_2,a_3,a_4}$ to the region $S_1 \subset R_1 \cap R_3$ gives a single-valued branch of the element on R_3, which is then called the *preferred branch on* R_3. Secondly, the restriction of the preferred branch on R_1 to the region $S_1 \subset R_1 \cap R_4$ also gives a single-valued branch of the element on R_4, which is then called the *preferred branch on* R_4. Moreover, the restriction of the preferred branch on R_4 to the region $S_2 \subset R_4 \cap R_2$ then gives a single-valued branch of the element on R_2 and we call it the *preferred branch on* R_2. It was verified in [H2, C] that $\mathbb{G}^{a_1,a_2,a_3,a_4}$ is a vector space.

For any element of $\mathbb{G}^{a_1,a_2,a_3,a_4}$, the preferred branches of this function on R_1, R_2 and R_3 give formal series in

$$\coprod_{a\in\mathcal{A}} x_1^{h_{a_4}-h_{a_1}-h_a} x_2^{h_a-h_{a_2}-h_{a_3}} \mathbb{C}[[x_2/x_1]][x_1, x_1^{-1}, x_2, x_2^{-1}], \tag{2.61}$$

$$\coprod_{a\in\mathcal{A}} x_2^{h_{a_4}-h_{a_2}-h_a} x_1^{h_a-h_{a_1}-h_{a_3}}\mathbb{C}[[x_1/x_2]][x_1,x_1^{-1},x_2,x_2^{-1}] \tag{2.62}$$

and

$$\coprod_{a\in\mathcal{A}} x_2^{h_{a_4}-h_a-h_{a_3}} x_0^{h_a-h_{a_1}-h_{a_2}}\mathbb{C}[[x_0/x_2]][x_0,x_0^{-1},x_2,x_2^{-1}], \tag{2.63}$$

respectively, which induce linear maps

$$\iota_{12}:\mathbb{G}^{a_1,a_2,a_3,a_4} \to \coprod_{a\in\mathcal{A}} x_1^{h_{a_4}-h_{a_1}-h_a} x_2^{h_a-h_{a_2}-h_{a_3}}\mathbb{C}[[x_2/x_1]][x_1,x_1^{-1},x_2,x_2^{-1}], \tag{2.64}$$

$$\iota_{21}:\mathbb{G}^{a_1,a_2,a_3,a_4} \to \coprod_{a\in\mathcal{A}} x_2^{h_{a_4}-h_{a_2}-h_a} x_1^{h_a-h_{a_1}-h_{a_3}}\mathbb{C}[[x_1/x_2]][x_1,x_1^{-1},x_2,x_2^{-1}], \tag{2.65}$$

$$\iota_{20}:\mathbb{G}^{a_1,a_2,a_3,a_4} \to \coprod_{a\in\mathcal{A}} x_2^{h_{a_4}-h_a-h_{a_3}} x_0^{h_a-h_{a_1}-h_{a_2}}\mathbb{C}[[x_0/x_2]][x_0,x_0^{-1},x_2,x_2^{-1}] \tag{2.66}$$

generalizing ι_{12}, ι_{21} and ι_{20} discussed at the beginning of this section. These maps are injective because analytic extensions are unique.

For $a_1,a_2,a_3,a_4\in\mathcal{A}$, $\mathbb{G}^{a_1,a_2,a_3,a_4}$ is a module over the ring $\mathbb{C}[x_1,x_1^{-1},x_2,x_2^{-1},(x_1-x_2)^{-1}]$. Huang [H2] proved the following lemma:

Lemma 2.6. *For any $a_1,a_2,a_3,a_4\in\mathcal{A}$, the module $\mathbb{G}^{a_1,a_2,a_3,a_4}$ is free.*

Remark 2.7. *In the following theorem for Jacobi identity and for the rest of the paper, we fix a basis $\{e_\alpha^{a_1,a_2,a_3,a_4}\}_{\alpha\in\mathbb{A}(a_1,a_2,a_3,a_4)}$ of the free module $\mathbb{G}^{a_1,a_2,a_3,a_4}$ over the ring $\mathbb{C}[x_1,x_1^{-1},x_2,x_2^{-1},(x_1-x_2)^{-1}]$ for any $a_1,a_2,a_3,a_4\in\mathcal{A}$, where $\mathbb{A}(a_1,a_2,a_3,a_4)$ is the index set of the basis.*

Now we give the Jacobi identity derived in [H2]:

Theorem 2.8 (Jacobi identity). *For any $a_1,a_2,a_3,a_4\in\mathcal{A}$, there exist linear maps*

$$\begin{aligned} f_\alpha^{a_1,a_2,a_3,a_4} : W^{a_1}\otimes W^{a_2}\otimes W^{a_3}\otimes \pi_P(\coprod_{a_5\in\mathcal{A}}\mathcal{V}_{a_1a_5}^{a_4}\otimes\mathcal{V}_{a_2a_3}^{a_5}) &\\ \to W^{a_4}[[x_2/x_1]][x_1,x_1^{-1},x_2,x_2^{-1}]&\\ w_{(a_1)}\otimes w_{(a_2)}\otimes w_{(a_3)}\otimes[\mathcal{Z}]_P &\\ \mapsto f_\alpha^{a_1,a_2,a_3,a_4}(w_{(a_1)},w_{(a_2)},w_{(a_3)},[\mathcal{Z}]_P;x_1,x_2),& \end{aligned} \tag{2.67}$$

$$\begin{aligned} g_\alpha^{a_1,a_2,a_3,a_4} : W^{a_1}\otimes W^{a_2}\otimes W^{a_3}\otimes \pi_P(\coprod_{a_5\in\mathcal{A}}\mathcal{V}_{a_2a_5}^{a_4}\otimes\mathcal{V}_{a_1a_3}^{a_5}) &\\ \to W^{a_4}[[x_1/x_2]][x_1,x_1^{-1},x_2,x_2^{-1}]&\\ w_{(a_1)}\otimes w_{(a_2)}\otimes w_{(a_3)}\otimes[\mathcal{Z}]_P &\\ \mapsto g_\alpha^{a_1,a_2,a_3,a_4}(w_{(a_1)},w_{(a_2)},w_{(a_3)},[\mathcal{Z}]_P;x_1,x_2)& \end{aligned} \tag{2.68}$$

and

$$\begin{aligned} h_{\alpha}^{a_1,a_2,a_3,a_4} : W^{a_1} \otimes W^{a_2} \otimes W^{a_3} \otimes \pi_I(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_2}^{a_5} \otimes \mathcal{V}_{a_5a_3}^{a_4}) \\ \to W^{a_4}[[x_0/x_2]][x_0, x_0^{-1}, x_2, x_2^{-1}] \\ w_{(a_1)} \otimes w_{(a_2)} \otimes w_{(a_3)} \otimes [\mathcal{Z}]_I \\ \mapsto h_{\alpha}^{a_1,a_2,a_3,a_4}(w_{(a_1)}, w_{(a_2)}, w_{(a_3)}, [\mathcal{Z}]_I; x_0, x_2) \end{aligned} \tag{2.69}$$

for $\alpha \in \mathbb{A}(a_1, a_2, a_3, a_4)$*, such that for any* $w_{(a_1)} \in W^{a_1}$*,* $w_{(a_2)} \in W^{a_2}$*,* $w_{(a_3)} \in W^{a_3}$*, and any*

$$\mathcal{Z} \in \coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5} \subset \coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5}, \tag{2.70}$$

only finitely many of

$$f_{\alpha}^{a_1,a_2,a_3,a_4}(w_{(a_1)}, w_{(a_2)}, w_{(a_3)}, [\mathcal{Z}]_P; x_1, x_2), \tag{2.71}$$

$$g_{\alpha}^{a_1,a_2,a_3,a_4}(w_{(a_1)}, w_{(a_2)}, w_{(a_3)}, \mathcal{B}([\mathcal{Z}]_P); x_1, x_2), \tag{2.72}$$

and

$$h_{\alpha}^{a_1,a_2,a_3,a_4}(w_{(a_1)}, w_{(a_2)}, w_{(a_3)}, \mathcal{F}([\mathcal{Z}]_P); x_0, x_2), \tag{2.73}$$

$\alpha \in \mathbb{A}(a_1, a_2, a_3, a_4)$*, are nonzero,*

$$\begin{aligned} &(\tilde{\mathbf{P}}([\mathcal{Z}]_P))(w_{(a_1)}, w_{(a_2)}, w_{(a_3)}; x_1, x_2) \\ &\quad = \sum_{\alpha\in\mathbb{A}(a_1,a_2,a_3,a_4)} f_{\alpha}^{a_1,a_2,a_3,a_4}(w_{(a_1)}, w_{(a_2)}, w_{(a_3)}, [\mathcal{Z}]_P; x_1, x_2)\iota_{12}\left(e_{\alpha}^{a_1,a_2,a_3,a_4}\right), \end{aligned} \tag{2.74}$$

$$\begin{aligned} &(\tilde{\mathbf{P}}(\mathcal{B}([\mathcal{Z}]_P)))(w_{(a_2)}, w_{(a_1)}, w_{(a_3)}; x_2, x_1) \\ &\quad = \sum_{\alpha\in\mathbb{A}(a_1,a_2,a_3,a_4)} g_{\alpha}^{a_1,a_2,a_3,a_4}(w_{(a_1)}, w_{(a_2)}, w_{(a_3)}, \mathcal{B}([\mathcal{Z}]_P); x_1, x_2)\iota_{21}\left(e_{\alpha}^{a_1,a_2,a_3,a_4}\right), \end{aligned} \tag{2.75}$$

$$\begin{aligned} &(\tilde{\mathbf{I}}(\mathcal{F}([\mathcal{Z}]_P)))(w_{(a_1)}, w_{(a_2)}, w_{(a_3)}; x_0, x_2) \\ &\quad = \sum_{\alpha\in\mathbb{A}(a_1,a_2,a_3,a_4)} h_{\alpha}^{a_1,a_2,a_3,a_4}(w_{(a_1)}, w_{(a_2)}, w_{(a_3)}, \mathcal{F}([\mathcal{Z}]_P); x_0, x_2)\iota_{20}(e_{\alpha}^{a_1,a_2,a_3,a_4}), \end{aligned} \tag{2.76}$$

and the following Jacobi identity holds:

$$\begin{aligned} &x_0^{-1}\delta\left(\frac{x_1 - x_2}{x_0}\right) f_{\alpha}^{a_1,a_2,a_3,a_4}(w_{(a_1)}, w_{(a_2)}, w_{(a_3)}, [\mathcal{Z}]_P; x_1, x_2) \\ &\qquad -x_0^{-1}\delta\left(\frac{x_2 - x_1}{-x_0}\right) g_{\alpha}^{a_1,a_2,a_3,a_4}(w_{(a_1)}, w_{(a_2)}, w_{(a_3)}, \mathcal{B}([\mathcal{Z}]_P); x_1, x_2) \\ &\quad = x_2^{-1}\delta\left(\frac{x_1 - x_0}{x_2}\right) h_{\alpha}^{a_1,a_2,a_3,a_4}(w_{(a_1)}, w_{(a_2)}, w_{(a_3)}, \mathcal{F}([\mathcal{Z}]_P); x_0, x_2) \end{aligned} \tag{2.77}$$

for $\alpha \in \mathbb{A}(a_1, a_2, a_3, a_4)$*.*

3 S_3-symmetry of the Jacobi identity

In this section, we formulate and prove a symmetric property of the Jacobi identity for intertwining operator algebras under the symmetric group S_3, which is the main result of this paper. Here is the precise statement of the main result:

Theorem 3.1. *In the presence of the axioms for an intertwining operator algebra except for the associativity property, we assume that there exists an isomorphism*

$$\mathcal{F}: \quad \frac{\coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4}\otimes\mathcal{V}_{a_2a_3}^{a_5}}{\text{Ker }\mathbf{P}} \longrightarrow \frac{\coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_2}^{a_5}\otimes\mathcal{V}_{a_5a_3}^{a_4}}{\text{Ker }\mathbf{I}} \tag{3.1}$$

satisfying

$$\mathcal{F}\left(\pi_P\left(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4}\otimes\mathcal{V}_{a_2a_3}^{a_5}\right)\right) = \pi_I\left(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_2}^{a_5}\otimes\mathcal{V}_{a_5a_3}^{a_4}\right) \tag{3.2}$$

for any $a_1,\cdots,a_4\in\mathcal{A}$, and that the Moore-Seiberg equations (2.50) and (2.51) hold, then the Jacobi identity for the ordered triple $(\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)})\in\coprod_{i=1}^3 W^{a_i}$ implies the Jacobi identity for the triple $(\tilde{w}_{(a_{\tau(1)})},\tilde{w}_{(a_{\tau(2)})},\tilde{w}_{(a_{\tau(3)})})\in\coprod_{i=1}^3 W^{a_{\tau(i)}}$ for any $\tau\in S_3$.

Remark 3.2. *In the above theorem, since there's no associativity in the assumptions, we have no fusing isomorphism. The assumption that the Moore-Seiberg equations (2.50) and (2.51) hold in fact means that they hold with the fusing isomorphism replaced by the given isomorphism $\mathcal{F}$ in (3.1). Moreover, from $\mathcal{F}$ in (3.1) and the isomorphism $\tilde{\Omega}^{(1)}$ we obtain an isomorphism*

$$\mathcal{B}=\mathcal{F}^{-1}\circ\tilde{\Omega}^{(1)}\circ\mathcal{F}: \quad \frac{\coprod\limits_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4}\otimes\mathcal{V}_{a_2a_3}^{a_5}}{\text{Ker }\mathbf{P}} \longrightarrow \frac{\coprod\limits_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_2a_5}^{a_4}\otimes\mathcal{V}_{a_1a_3}^{a_5}}{\text{Ker }\mathbf{P}}. \tag{3.3}$$

And we also assume that the isomorphisms $\mathcal{F}$ and $\mathcal{B}$ involved in the formulation of the Jacobi identity are replaced by the given isomorphism in (3.1) and the deduced isomorphism in (3.3), respectively.

The rest of this section is devoted to proving the S_3-symmetry of the Jacobi identity. We achieve this goal by establishing three results that lead to Theorem 3.1.

Proposition 3.3. *In the presence of the axioms for an intertwining operator algebra except for the associativity property, we assume that there exists an isomorphism*

$$\mathcal{F}: \quad \frac{\coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4}\otimes\mathcal{V}_{a_2a_3}^{a_5}}{\text{Ker }\mathbf{P}} \longrightarrow \frac{\coprod_{a_1,a_2,a_3,a_4,a_5\in\mathcal{A}} \mathcal{V}_{a_1a_2}^{a_5}\otimes\mathcal{V}_{a_5a_3}^{a_4}}{\text{Ker }\mathbf{I}} \tag{3.4}$$

satisfying

$$\mathcal{F}\left(\pi_P\left(\coprod_{a_5\in\mathcal{A}}\mathcal{V}_{a_1a_5}^{a_4}\otimes\mathcal{V}_{a_2a_3}^{a_5}\right)\right)=\pi_I\left(\coprod_{a_5\in\mathcal{A}}\mathcal{V}_{a_1a_2}^{a_5}\otimes\mathcal{V}_{a_5a_3}^{a_4}\right) \tag{3.5}$$

for any $a_1,\cdots,a_4\in\mathcal{A}$, *and that the Jacobi identity for the ordered triple* $(\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)})\in\coprod_{i=1}^3 W^{a_i}$ *holds, then for any* $a_4\in\mathcal{A}$, $w'_{(a_4)}\in(W^{a_4})'$ *and* $\mathcal{Z}\in\coprod_{a_5\in\mathcal{A}}\mathcal{V}_{a_1a_5}^{a_4}\otimes\mathcal{V}_{a_2a_3}^{a_5}$, *there exists a multivalued analytic function*

$$\Phi(w'_{(a_4)},\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)},[\mathcal{Z}]_P;z_1,z_2)\in\mathbb{G}^{a_1,a_2,a_3,a_4} \tag{3.6}$$

such that

$$\langle w'_{(a_4)},(\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)};x_1,x_2)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1},x_2^n=e^{n\log z_2}}, \tag{3.7}$$

$$\langle w'_{(a_4)},(\tilde{\mathbf{P}}(\mathcal{B}([\mathcal{Z}]_P)))(\tilde{w}_{(a_2)},\tilde{w}_{(a_1)},\tilde{w}_{(a_3)};x_2,x_1)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1},x_2^n=e^{n\log z_2}}, \tag{3.8}$$

$$\langle w'_{(a_4)},(\tilde{\mathbf{I}}(\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)};x_0,x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)},x_2^n=e^{n\log z_2}} \tag{3.9}$$

and

$$\langle w'_{(a_4)},(\tilde{\mathbf{I}}(\tilde{\Omega}^{(1)}(\mathcal{F}([\mathcal{Z}]_P))))(\tilde{w}_{(a_2)},\tilde{w}_{(a_1)},\tilde{w}_{(a_3)};x_0,x_1)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_2-z_1)},x_1^n=e^{n\log z_1}} \tag{3.10}$$

are its preferred branches on R_1, R_2, R_3 *and* R_4, *respectively. Moreover,*

$$\begin{aligned}&\langle w'_{(a_4)},(\tilde{\mathbf{I}}(\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)};x_0,x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)},x_2^n=e^{n\log z_2}}\\&\quad=\langle w'_{(a_4)},(\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)};x_1,x_2)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1},x_2^n=e^{n\log z_2}}\end{aligned} \tag{3.11}$$

on the region

$$S_1=\{(z_1,z_2)\in\mathbb{C}^2\mid \mathrm{Re}z_1>\mathrm{Re}z_2>\mathrm{Re}(z_1-z_2)>0,\ \mathrm{Im}z_1>\mathrm{Im}z_2>\mathrm{Im}(z_1-z_2)>0\},$$

and

$$\begin{aligned}&\langle w'_{(a_4)},(\tilde{\mathbf{I}}(\tilde{\Omega}^{(1)}(\mathcal{F}([\mathcal{Z}]_P))))(\tilde{w}_{(a_2)},\tilde{w}_{(a_1)},\tilde{w}_{(a_3)};x_0,x_1)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_2-z_1)},x_1^n=e^{n\log z_1}}\\&\quad=\langle w'_{(a_4)},(\tilde{\mathbf{P}}(\mathcal{B}([\mathcal{Z}]_P)))(\tilde{w}_{(a_2)},\tilde{w}_{(a_1)},\tilde{w}_{(a_3)};x_2,x_1)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1},x_2^n=e^{n\log z_2}}\end{aligned} \tag{3.12}$$

on the region

$$S_2=\{(z_1,z_2)\in\mathbb{C}^2\mid \mathrm{Re}z_2>\mathrm{Re}z_1>\mathrm{Re}(z_2-z_1)>0,\ \mathrm{Im}z_2>\mathrm{Im}z_1>\mathrm{Im}(z_2-z_1)>0\}.$$

Remark 3.4. *In the above proposition, the formulations involving the isomorphisms* $\mathcal{F}$ *and* $\mathcal{B}$ *have the same assumptions as we discussed in Remark 3.2 below Theorem 3.1.*

Remark 3.5. *In [C], we proved that for intertwining operator algebras, the generalized rationality, commutativity and associativity follow from the Jacobi identity. And its proof involves only one ordered triple* $(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}) \in \coprod_{i=1}^{3} W^{a_i}$ *for both commutativity, associativity and the Jacobi identity. Minus the skew-symmetry condition, the above proposition becomes a one-ordered-triple version of Theorem 3.3 in [C]. However, for the sake of proving Theorem 3.1, we add the extra skew-symmetry condition in the above proposition to obtain the preferred branch (3.10) on* R_4 *and the analytic extension relation (3.12).*

Proof of Proposition 3.3. Since the Jacobi identity holds for the ordered triple $(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}) \in \coprod_{i=1}^{3} W^{a_i}$, then for any $a_4 \in \mathcal{A}$, there exist linear maps

$$\begin{aligned} f_{\alpha}^{a_1,a_2,a_3,a_4} : W^{a_1} \otimes W^{a_2} \otimes W^{a_3} \otimes \pi_P(\coprod_{a_5 \in \mathcal{A}} \mathcal{V}_{a_1 a_5}^{a_4} \otimes \mathcal{V}_{a_2 a_3}^{a_5}) & \\ \to W^{a_4}[[x_2/x_1]][x_1, x_1^{-1}, x_2, x_2^{-1}] & \\ w_{(a_1)} \otimes w_{(a_2)} \otimes w_{(a_3)} \otimes [\mathcal{Z}]_P & \\ \mapsto f_{\alpha}^{a_1,a_2,a_3,a_4}(w_{(a_1)}, w_{(a_2)}, w_{(a_3)}, [\mathcal{Z}]_P; x_1, x_2), & \end{aligned} \tag{3.13}$$

$$\begin{aligned} g_{\alpha}^{a_1,a_2,a_3,a_4} : W^{a_1} \otimes W^{a_2} \otimes W^{a_3} \otimes \pi_P(\coprod_{a_5 \in \mathcal{A}} \mathcal{V}_{a_2 a_5}^{a_4} \otimes \mathcal{V}_{a_1 a_3}^{a_5}) & \\ \to W^{a_4}[[x_1/x_2]][x_1, x_1^{-1}, x_2, x_2^{-1}] & \\ w_{(a_1)} \otimes w_{(a_2)} \otimes w_{(a_3)} \otimes [\mathcal{Z}]_P & \\ \mapsto g_{\alpha}^{a_1,a_2,a_3,a_4}(w_{(a_1)}, w_{(a_2)}, w_{(a_3)}, [\mathcal{Z}]_P; x_1, x_2) & \end{aligned} \tag{3.14}$$

and

$$\begin{aligned} h_{\alpha}^{a_1,a_2,a_3,a_4} : W^{a_1} \otimes W^{a_2} \otimes W^{a_3} \otimes \pi_I(\coprod_{a_5 \in \mathcal{A}} \mathcal{V}_{a_1 a_2}^{a_5} \otimes \mathcal{V}_{a_5 a_3}^{a_4}) & \\ \to W^{a_4}[[x_0/x_2]][x_0, x_0^{-1}, x_2, x_2^{-1}] & \\ w_{(a_1)} \otimes w_{(a_2)} \otimes w_{(a_3)} \otimes [\mathcal{Z}]_I & \\ \mapsto h_{\alpha}^{a_1,a_2,a_3,a_4}(w_{(a_1)}, w_{(a_2)}, w_{(a_3)}, [\mathcal{Z}]_I; x_0, x_2) & \end{aligned} \tag{3.15}$$

for $\alpha \in \mathbb{A}(a_1, a_2, a_3, a_4)$, such that for any

$$\mathcal{Z} \in \coprod_{a_5 \in \mathcal{A}} \mathcal{V}_{a_1 a_5}^{a_4} \otimes \mathcal{V}_{a_2 a_3}^{a_5} \subset \coprod_{a_1,a_2,a_3,a_4,a_5 \in \mathcal{A}} \mathcal{V}_{a_1 a_5}^{a_4} \otimes \mathcal{V}_{a_2 a_3}^{a_5}, \tag{3.16}$$

only finitely many of

$$f_{\alpha}^{a_1,a_2,a_3,a_4}(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; x_1, x_2), \tag{3.17}$$

$$g_{\alpha}^{a_1,a_2,a_3,a_4}(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, \mathcal{B}([\mathcal{Z}]_P); x_1, x_2), \tag{3.18}$$

and

$$h_{\alpha}^{a_1,a_2,a_3,a_4}(\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)},\mathcal{F}([\mathcal{Z}]_P);x_0,x_2), \tag{3.19}$$

$\alpha\in\mathbb{A}(a_1,a_2,a_3,a_4)$, are nonzero,

$$\begin{aligned}&(\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)};x_1,x_2)\\&\quad=\sum_{\alpha\in\mathbb{A}(a_1,a_2,a_3,a_4)} f_{\alpha}^{a_1,a_2,a_3,a_4}(\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)},[\mathcal{Z}]_P;x_1,x_2)\iota_{12}\left(e_{\alpha}^{a_1,a_2,a_3,a_4}\right),\end{aligned} \tag{3.20}$$

$$\begin{aligned}&(\tilde{\mathbf{P}}(\mathcal{B}([\mathcal{Z}]_P)))(\tilde{w}_{(a_2)},\tilde{w}_{(a_1)},\tilde{w}_{(a_3)};x_2,x_1)\\&\quad=\sum_{\alpha\in\mathbb{A}(a_1,a_2,a_3,a_4)} g_{\alpha}^{a_1,a_2,a_3,a_4}(\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)},\mathcal{B}([\mathcal{Z}]_P);x_1,x_2)\iota_{21}\left(e_{\alpha}^{a_1,a_2,a_3,a_4}\right),\end{aligned} \tag{3.21}$$

$$\begin{aligned}&(\tilde{\mathbf{I}}(\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)};x_0,x_2)\\&\quad=\sum_{\alpha\in\mathbb{A}(a_1,a_2,a_3,a_4)} h_{\alpha}^{a_1,a_2,a_3,a_4}(\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)},\mathcal{F}([\mathcal{Z}]_P);x_0,x_2)\iota_{20}(e_{\alpha}^{a_1,a_2,a_3,a_4}),\end{aligned} \tag{3.22}$$

and the following *Jacobi identity* holds:

$$\begin{aligned}&x_0^{-1}\delta\left(\frac{x_1-x_2}{x_0}\right)f_{\alpha}^{a_1,a_2,a_3,a_4}(\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)},[\mathcal{Z}]_P;x_1,x_2)\\&\quad-x_0^{-1}\delta\left(\frac{x_2-x_1}{-x_0}\right)g_{\alpha}^{a_1,a_2,a_3,a_4}(\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)},\mathcal{B}([\mathcal{Z}]_P);x_1,x_2)\\&\quad=x_2^{-1}\delta\left(\frac{x_1-x_0}{x_2}\right)h_{\alpha}^{a_1,a_2,a_3,a_4}(\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)},\mathcal{F}([\mathcal{Z}]_P);x_0,x_2)\end{aligned} \tag{3.23}$$

for $\alpha\in\mathbb{A}(a_1,a_2,a_3,a_4)$.

In analogy with the proof of Theorem 3.3 in [C], we can obtain that, for any $a_4\in\mathcal{A}$, $\alpha\in\mathbb{A}(a_1,a_2,a_3,a_4)$, there exists linear map

$$\begin{aligned}F_{\alpha}:\quad&(W^{a_4})'\otimes W^{a_1}\otimes W^{a_2}\otimes W^{a_3}\otimes\pi_P(\coprod_{a_5\in\mathcal{A}}\mathcal{V}_{a_1a_5}^{a_4}\otimes\mathcal{V}_{a_2a_3}^{a_5})\\&\longrightarrow\mathbb{C}[x_1,x_1^{-1},x_2,x_2^{-1},(x_1-x_2)^{-1}]\end{aligned} \tag{3.24}$$

with $F_{\alpha}(w'_{(a_4)},\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)},[\mathcal{Z}]_P;x_1,x_2)$ denoted by the image of

$$w'_{(a_4)}\otimes\tilde{w}_{(a_1)}\otimes\tilde{w}_{(a_2)}\otimes\tilde{w}_{(a_3)}\otimes[\mathcal{Z}]_P$$

under F_{α}, such that

$$\begin{aligned}&\langle w'_{(a_4)},f_{\alpha}^{a_1,a_2,a_3,a_4}(\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)},[\mathcal{Z}]_P;x_1,x_2)\rangle_{W^{a_4}}\\&\quad=\iota_{12}F_{\alpha}(w'_{(a_4)},\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)},[\mathcal{Z}]_P;x_1,x_2),\end{aligned} \tag{3.25}$$

$$\begin{aligned}\langle w'_{(a_4)}, g_\alpha^{a_1,a_2,a_3,a_4}(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, \mathcal{B}([\mathcal{Z}]_P); x_1, x_2)\rangle_{W^{a_4}} \\ = \iota_{21} F_\alpha(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; x_1, x_2)\end{aligned} \tag{3.26}$$

and

$$\begin{aligned}\langle w'_{(a_4)}, h_\alpha^{a_1,a_2,a_3,a_4}(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, \mathcal{F}([\mathcal{Z}]_P); x_0, x_2)\rangle_{W^{a_4}} \\ = \iota_{20} F_\alpha(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; x_2 + x_0, x_2)\end{aligned} \tag{3.27}$$

for $\alpha \in \mathbb{A}(a_1, a_2, a_3, a_4)$. Moreover, since $\mathbb{G}^{a_1,a_2,a_3,a_4}$ is a free module over the ring

$$\mathbb{C}[x_1, x_1^{-1}, x_2, x_2^{-1}, (x_1 - x_2)^{-1}] \tag{3.28}$$

with a basis $\{e_\alpha^{a_1,a_2,a_3,a_4}\}_{\alpha\in\mathbb{A}(a_1,a_2,a_3,a_4)}$, we have

$$\begin{aligned}\mathbb{G}^{a_1,a_2,a_3,a_4} \ni\ & \Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1, z_2) \\ = & \sum_{\alpha\in\mathbb{A}(a_1,a_2,a_3,a_4)} F_\alpha(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; x_1, x_2) e_\alpha^{a_1,a_2,a_3,a_4},\end{aligned} \tag{3.29}$$

and

$$\begin{aligned}\langle w'_{(a_4)}, (\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_1, x_2)\rangle_{W^{a_4}} \\ = \iota_{12} \Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1, z_2),\end{aligned} \tag{3.30}$$

$$\begin{aligned}\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}([\mathcal{Z}]_P)))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_2, x_1)\rangle_{W^{a_4}} \\ = \iota_{21} \Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1, z_2),\end{aligned} \tag{3.31}$$

$$\begin{aligned}\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}} \\ = \iota_{20} \Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1, z_2).\end{aligned} \tag{3.32}$$

So the preferred branches of $\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1, z_2)$ on R_1, R_2 and R_3 are

$$\langle w'_{(a_4)}, (\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}}, \tag{3.33}$$

$$\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}([\mathcal{Z}]_P)))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_2, x_1)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}} \tag{3.34}$$

and

$$\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)}, x_2^n=e^{n\log z_2}}, \tag{3.35}$$

respectively. And the multivalued analytic functions

$$\langle w'_{(a_4)}, (\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1=z_1, x_2=z_2}, \tag{3.36}$$

$$\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}([\mathcal{Z}]_P)))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_2, x_1)\rangle_{W^{a_4}}\big|_{x_1=z_1, x_2=z_2} \tag{3.37}$$

and

$$\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0=z_1-z_2, x_2=z_2} \tag{3.38}$$

are restrictions of the multivalued analytic function $\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1, z_2)$ to their domains $|z_1| > |z_2| > 0$, $|z_2| > |z_1| > 0$ and $|z_2| > |z_1 - z_2| > 0$ respectively.

Then by the definition of the preferred branch of an element of $\mathbb{G}^{a_1,a_2,a_3,a_4}$ on R_3, we can deduce that

$$\begin{aligned}
&\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)}, x_2^n=e^{n\log z_2}}\\
&\quad = \langle w'_{(a_4)}, (\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}}
\end{aligned} \tag{3.39}$$

on the region

$$S_1 = \{(z_1, z_2) \in \mathbb{C}^2 \mid \mathrm{Re} z_1 > \mathrm{Re} z_2 > \mathrm{Re}(z_1 - z_2) > 0,\ \mathrm{Im} z_1 > \mathrm{Im} z_2 > \mathrm{Im}(z_1 - z_2) > 0\}.$$

Moreover, by skew-symmetry and (3.39), we see that

$$\begin{aligned}
&\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\tilde{\Omega}^{(1)}(\mathcal{F}([\mathcal{Z}]_P))))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_0, x_1)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_2-z_1)}, x_1^n=e^{n\log z_1}}\\
&\quad = \langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; e^{-\pi i}x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_2-z_1)}, x_2^n=e^{n\log z_2}}\\
&\quad = \langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)}, x_2^n=e^{n\log z_2}}\\
&\quad = \langle w'_{(a_4)}, (\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}}
\end{aligned} \tag{3.40}$$

on the region S_1. So by the definition of the preferred branch of an element of $\mathbb{G}^{a_1,a_2,a_3,a_4}$ on R_4, we deduce that the preferred branch on R_4 of $\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1, z_2)$ is equal to the single-valued analytic function

$$\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\tilde{\Omega}^{(1)}(\mathcal{F}([\mathcal{Z}]_P))))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_0, x_1)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_2-z_1)}, x_1^n=e^{n\log z_1}} \tag{3.41}$$

on the region S_1. Since the preferred branch on R_4 of $\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1, z_2)$ and the function (3.41) are both single-valued analytic functions on the domain R_4 which contains S_1, by the basic properties of analytic functions we conclude that they are equal on R_4; namely, the single-valued analytic function (3.41) defined on the region R_4 is the preferred branch of $\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1, z_2)$ on R_4. Furthermore, by the definition of the preferred branch of an element of $\mathbb{G}^{a_1,a_2,a_3,a_4}$ on R_2, we can conclude that

$$\begin{aligned}
&\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\tilde{\Omega}^{(1)}(\mathcal{F}([\mathcal{Z}]_P))))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_0, x_1)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_2-z_1)}, x_1^n=e^{n\log z_1}}\\
&\quad = \langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}([\mathcal{Z}]_P)))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_2, x_1)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}}
\end{aligned} \tag{3.42}$$

on the region

$$S_2 = \{(z_1, z_2) \in \mathbb{C}^2 \mid \mathrm{Re} z_2 > \mathrm{Re} z_1 > \mathrm{Re}(z_2 - z_1) > 0,\ \mathrm{Im} z_2 > \mathrm{Im} z_1 > \mathrm{Im}(z_2 - z_1) > 0\}.$$

So this proposition holds. ∎

Theorem 3.6. *Assume that the assumptions of Theorem 3.1 hold, then the Jacobi identity holds for the ordered triple* $(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)})$.

Proof. Consider any $a_4 \in \mathcal{A}$, $w'_{(a_4)} \in (W^{a_4})'$ and $\mathcal{Z} \in \coprod_{a_5 \in \mathcal{A}} \mathcal{V}_{a_1 a_5}^{a_4} \otimes \mathcal{V}_{a_2 a_3}^{a_5}$. Since the assumptions of Theorem 3.1 contain the assumptions of Proposition 3.3, we obtain a multivalued analytic function $\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1, z_2)$ (see (3.6)) such that (3.7)-(3.10) are its preferred branches on R_1, R_2, R_3 and R_4, respectively. Moreover, the preferred branches on R_1, R_2, R_3 and R_4 have relations (3.11)-(3.12). Interchanging z_1 and z_2 in the multivalued analytic function $\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1, z_2)$, we obtain another multivalued analytic function

$$\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_2, z_1) \tag{3.43}$$

on $M^2 = \{(z_1, z_2) \in \mathbb{C}^2 \mid z_1, z_2 \neq 0, z_1 \neq z_2\}$. By interchanging z_1 and z_2 in (3.8), we see that

$$\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}([\mathcal{Z}]_P)))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1^n = e^{n \log z_1}, x_2^n = e^{n \log z_2}} \tag{3.44}$$

is a branch of $\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_2, z_1)$ on the region R_1.

Moreover, by interchanging z_1 and z_2 in (3.12), we get

$$\begin{aligned}
&\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\tilde{\Omega}^{(1)}(\mathcal{F}([\mathcal{Z}]_P))))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n = e^{n \log(z_1 - z_2)}, x_2^n = e^{n \log z_2}} \\
&\quad = \langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}([\mathcal{Z}]_P)))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1^n = e^{n \log z_1}, x_2^n = e^{n \log z_2}}
\end{aligned} \tag{3.45}$$

on the region S_1. Since the single-valued analytic function

$$\begin{aligned}
&\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}(\mathcal{B}([\mathcal{Z}]_P))))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n = e^{n \log(z_1 - z_2)}, x_2^n = e^{n \log z_2}} \\
&\quad = \langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\tilde{\Omega}^{(1)}(\mathcal{F}([\mathcal{Z}]_P))))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n = e^{n \log(z_1 - z_2)}, x_2^n = e^{n \log z_2}}
\end{aligned} \tag{3.46}$$

on S_1 can be naturally analytically extended to the region R_3, we therefore get a branch of $\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_2, z_1)$ on the region R_3.

By (3.45) and skew-symmetry, we have

$$\begin{aligned}
&\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\tilde{\Omega}^{(1)}\tilde{\Omega}^{(1)}(\mathcal{F}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_1)\rangle_{W^{a_4}}\big|_{x_0^n = e^{n \log(z_2 - z_1)}, x_1^n = e^{n \log z_1}} \\
&\quad = \langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\tilde{\Omega}^{(1)}(\mathcal{F}([\mathcal{Z}]_P))))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; e^{-\pi i} x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n = e^{n \log(z_2 - z_1)}, x_2^n = e^{n \log z_2}} \\
&\quad = \langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\tilde{\Omega}^{(1)}(\mathcal{F}([\mathcal{Z}]_P))))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n = e^{n \log(z_1 - z_2)}, x_2^n = e^{n \log z_2}} \\
&\quad = \langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}([\mathcal{Z}]_P)))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1^n = e^{n \log z_1}, x_2^n = e^{n \log z_2}}
\end{aligned} \tag{3.47}$$

on the region S_1. Moreover, the first line of (3.47) on S_1 can be naturally analytically extended to the region R_4. So the single-valued analytic function

$$\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\tilde{\Omega}^{(1)}\tilde{\Omega}^{(1)}(\mathcal{F}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_1)\rangle_{W^{a_4}}|_{x_0^n=e^{n\log(z_2-z_1)}, x_1^n=e^{n\log z_1}} \tag{3.48}$$

defined on the region R_4 is a branch of $\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_2, z_1)$ on the region R_4.

Observe that $[\mathcal{Z}]_P \in \pi_P(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5})$ implies

$$\mathcal{B}(\mathcal{B}([\mathcal{Z}]_P)) \in \pi_P(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5}). \tag{3.49}$$

And since (3.6)-(3.12) hold for any $[\mathcal{Z}]_P \in \pi_P(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5})$, they should hold with $[\mathcal{Z}]_P$ replaced by $\mathcal{B}(\mathcal{B}([\mathcal{Z}]_P))$ for any $[\mathcal{Z}]_P \in \pi_P(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5})$. So replacing $[\mathcal{Z}]_P$ by $\mathcal{B}(\mathcal{B}([\mathcal{Z}]_P))$ in (3.11), we get

$$\begin{aligned}&\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}(\mathcal{B}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_1, x_2)\rangle_{W^{a_4}}|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}}\\ &\quad= \langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}(\mathcal{B}(\mathcal{B}([\mathcal{Z}]_P)))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}|_{x_0^n=e^{n\log(z_1-z_2)}, x_2^n=e^{n\log z_2}}\end{aligned} \tag{3.50}$$

on the region S_1. Interchanging z_1 and z_2 in (3.50), we obtain

$$\begin{aligned}&\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}(\mathcal{B}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_2, x_1)\rangle_{W^{a_4}}|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}}\\ &\quad= \langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}(\mathcal{B}(\mathcal{B}([\mathcal{Z}]_P)))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_1)\rangle_{W^{a_4}}|_{x_0^n=e^{n\log(z_2-z_1)}, x_1^n=e^{n\log z_1}}\\ &\quad= \langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\tilde{\Omega}^{(1)}\tilde{\Omega}^{(1)}\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_1)\rangle_{W^{a_4}}|_{x_0^n=e^{n\log(z_2-z_1)}, x_1^n=e^{n\log z_1}}\end{aligned} \tag{3.51}$$

on the region S_2. Moreover, the first line of (3.51) on S_2 can be naturally analytically extended to the region R_2. So the single-valued analytic function

$$\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}(\mathcal{B}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_2, x_1)\rangle_{W^{a_4}}|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}} \tag{3.52}$$

defined on the region R_2 is a branch of $\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_2, z_1)$ on R_2.

From the above discussion (3.44)-(3.52), we see that the multivalued analytic functions

$$\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}([\mathcal{Z}]_P)))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_1, x_2)\rangle_{W^{a_4}}|_{x_1=z_1, x_2=z_2}, \tag{3.53}$$

$$\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}(\mathcal{B}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_2, x_1)\rangle_{W^{a_4}}|_{x_1=z_1, x_2=z_2} \tag{3.54}$$

and

$$\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}(\mathcal{B}([\mathcal{Z}]_P))))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}|_{x_0=z_1-z_2, x_2=z_2} \tag{3.55}$$

are restrictions of the multivalued analytic function $\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_2, z_1)$ to their domains $|z_1| > |z_2| > 0$, $|z_2| > |z_1| > 0$ and $|z_2| > |z_1 - z_2| > 0$ respectively. So with the branch (3.44) chosen as the preferred branch on R_1, $\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_2, z_1)$

becomes an element of $\mathbb{G}^{a_2,a_1,a_3,a_4}$. Moreover, by (3.45), (3.46), (3.47), (3.48), (3.51), (3.52), and by the definition of the preferred branches of an element of $\mathbb{G}^{a_2,a_1,a_3,a_4}$ on R_2 and R_3, we see that

$$\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}(\mathcal{B}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_2, x_1)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}} \tag{3.56}$$

and

$$\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}(\mathcal{B}([\mathcal{Z}]_P))))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)}, x_2^n=e^{n\log z_2}} \tag{3.57}$$

are the preferred branches of $\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_2, z_1)$ on R_2 and R_3 respectively. Therefore, we have

$$\begin{aligned}\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}([\mathcal{Z}]_P)))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_1, x_2)\rangle_{W^{a_4}}\\ = \iota_{12}\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_2, z_1),\end{aligned} \tag{3.58}$$

$$\begin{aligned}\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}(\mathcal{B}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_2, x_1)\rangle_{W^{a_4}}\\ = \iota_{21}\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_2, z_1),\end{aligned} \tag{3.59}$$

$$\begin{aligned}\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}(\mathcal{B}([\mathcal{Z}]_P))))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}\\ = \iota_{20}\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_2, z_1).\end{aligned} \tag{3.60}$$

Let $\{e_\alpha^{a_2,a_1,a_3,a_4}\}_{\alpha\in\mathbb{A}(a_2,a_1,a_3,a_4)}$ be a basis of $\mathbb{G}^{a_2,a_1,a_3,a_4}$ over the ring

$$\mathbb{C}[x_1, x_1^{-1}, x_2, x_2^{-1}, (x_1-x_2)^{-1}]. \tag{3.61}$$

Then there exists unique

$$G_\alpha(w'_{(a_4)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \mathcal{B}([\mathcal{Z}]_P); x_1, x_2) \in \mathbb{C}[x_1, x_1^{-1}, x_2, x_2^{-1}, (x_1-x_2)^{-1}] \tag{3.62}$$

for $\alpha\in\mathbb{A}(a_2,a_1,a_3,a_4)$, such that only finitely many of them are nonzero and

$$\begin{aligned}\Phi(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_2, z_1)\\ = \sum_{\alpha\in\mathbb{A}(a_2,a_1,a_3,a_4)} G_\alpha(w'_{(a_4)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \mathcal{B}([\mathcal{Z}]_P); x_1, x_2)e_\alpha^{a_2,a_1,a_3,a_4}.\end{aligned} \tag{3.63}$$

By (2.36) and (3.2), we see that $\mathcal{B} = \mathcal{F}^{-1}\tilde{\Omega}^{(1)}\mathcal{F}$ is an isomorphism and that

$$\mathcal{B}(\pi_P(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4}\otimes\mathcal{V}_{a_2a_3}^{a_5})) = \pi_P(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_2a_5}^{a_4}\otimes\mathcal{V}_{a_1a_3}^{a_5}) \tag{3.64}$$

for any $a_1, \cdots, a_4 \in \mathcal{A}$. So we can define linear maps

$$\begin{aligned} & f_\alpha^{a_2,a_1,a_3,a_4}(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}) : \\ & \qquad \pi_P(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_2a_5}^{a_4} \otimes \mathcal{V}_{a_1a_3}^{a_5}) \to W^{a_4}[x_1, x_1^{-1}, x_2, x_2^{-1}][[x_2/x_1]], \end{aligned} \tag{3.65}$$

$$\begin{aligned} & g_\alpha^{a_2,a_1,a_3,a_4}(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}) : \\ & \qquad \pi_P(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5}) \to W^{a_4}[x_1, x_1^{-1}, x_2, x_2^{-1}][[x_1/x_2]], \end{aligned} \tag{3.66}$$

$$\begin{aligned} & h_\alpha^{a_2,a_1,a_3,a_4}(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}) : \\ & \qquad \pi_I(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_2a_1}^{a_5} \otimes \mathcal{V}_{a_5a_3}^{a_4}) \to W^{a_4}[x_0, x_0^{-1}, x_2, x_2^{-1}][[x_0/x_2]] \end{aligned} \tag{3.67}$$

by

$$\begin{aligned} & \langle w'_{(a_4)}, f_\alpha^{a_2,a_1,a_3,a_4}(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \mathcal{B}([\mathcal{Z}]_P); x_1, x_2)\rangle_{W^{a_4}} \\ & \qquad = \iota_{12} G_\alpha(w'_{(a_4)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \mathcal{B}([\mathcal{Z}]_P); x_1, x_2), \end{aligned} \tag{3.68}$$

$$\begin{aligned} & \langle w'_{(a_4)}, g_\alpha^{a_2,a_1,a_3,a_4}(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \mathcal{B}(\mathcal{B}([\mathcal{Z}]_P)); x_1, x_2)\rangle_{W^{a_4}} \\ & \qquad = \iota_{21} G_\alpha(w'_{(a_4)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \mathcal{B}([\mathcal{Z}]_P); x_1, x_2), \end{aligned} \tag{3.69}$$

$$\begin{aligned} & \langle w'_{(a_4)}, h_\alpha^{a_2,a_1,a_3,a_4}(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \mathcal{F}(\mathcal{B}([\mathcal{Z}]_P)); x_0, x_2)\rangle_{W^{a_4}} \\ & \qquad = \iota_{20} G_\alpha(w'_{(a_4)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \mathcal{B}([\mathcal{Z}]_P); x_2 + x_0, x_2) \end{aligned} \tag{3.70}$$

for $w'_{(a_4)} \in (W^{a_4})'$, $\mathcal{Z} \in \coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5}$ and $\alpha \in \mathbb{A}(a_2, a_1, a_3, a_4)$. Then by (3.58)-(3.60) and (3.63), we have

$$\begin{aligned} & (\tilde{\mathbf{P}}(\mathcal{B}([\mathcal{Z}]_P)))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_1, x_2) \\ & \qquad = \sum_{\alpha\in\mathbb{A}(a_2,a_1,a_3,a_4)} f_\alpha^{a_2,a_1,a_3,a_4}(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \mathcal{B}([\mathcal{Z}]_P); x_1, x_2)\iota_{12}(e_\alpha^{a_2,a_1,a_3,a_4}), \end{aligned} \tag{3.71}$$

$$\begin{aligned} & (\tilde{\mathbf{P}}(\mathcal{B}(\mathcal{B}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_2, x_1) \\ & \qquad = \sum_{\alpha\in\mathbb{A}(a_2,a_1,a_3,a_4)} g_\alpha^{a_2,a_1,a_3,a_4}(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \mathcal{B}(\mathcal{B}([\mathcal{Z}]_P)); x_1, x_2)\iota_{21}(e_\alpha^{a_2,a_1,a_3,a_4}), \end{aligned} \tag{3.72}$$

$$\begin{aligned} & (\tilde{\mathbf{I}}(\mathcal{F}(\mathcal{B}([\mathcal{Z}]_P))))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_0, x_2) \\ & \qquad = \sum_{\alpha\in\mathbb{A}(a_2,a_1,a_3,a_4)} h_\alpha^{a_2,a_1,a_3,a_4}(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \mathcal{F}(\mathcal{B}([\mathcal{Z}]_P)); x_0, x_2)\iota_{20}(e_\alpha^{a_2,a_1,a_3,a_4}). \end{aligned} \tag{3.73}$$

Moreover, by (3.62) and Proposition 2.1, we have

$$\begin{aligned}
&x_0^{-1}\delta\left(\frac{x_1-x_2}{x_0}\right)\iota_{12}G_\alpha(w'_{(a_4)},\tilde{w}_{(a_2)},\tilde{w}_{(a_1)},\tilde{w}_{(a_3)},\mathcal{B}([\mathcal{Z}]_P);x_1,x_2)\\
&\quad -x_0^{-1}\delta\left(\frac{x_2-x_1}{-x_0}\right)\iota_{21}G_\alpha(w'_{(a_4)},\tilde{w}_{(a_2)},\tilde{w}_{(a_1)},\tilde{w}_{(a_3)},\mathcal{B}([\mathcal{Z}]_P);x_1,x_2)\\
&\quad = x_2^{-1}\delta\left(\frac{x_1-x_0}{x_2}\right)\iota_{20}G_\alpha(w'_{(a_4)},\tilde{w}_{(a_2)},\tilde{w}_{(a_1)},\tilde{w}_{(a_3)},\mathcal{B}([\mathcal{Z}]_P);x_2+x_0,x_2) \qquad (3.74)
\end{aligned}$$

for $\alpha\in\mathbb{A}(a_2,a_1,a_3,a_4)$. Since $w'_{(a_4)}\in(W^{a_4})'$ is arbitrary, by (3.68)-(3.70) and (3.74), the Jacobi identity holds for the ordered triple $(\tilde{w}_{(a_2)},\tilde{w}_{(a_1)},\tilde{w}_{(a_3)})$:

$$\begin{aligned}
&x_0^{-1}\delta\left(\frac{x_1-x_2}{x_0}\right)f_\alpha^{a_2,a_1,a_3,a_4}(\tilde{w}_{(a_2)},\tilde{w}_{(a_1)},\tilde{w}_{(a_3)},\mathcal{B}([\mathcal{Z}]_P);x_1,x_2)\\
&\quad -x_0^{-1}\delta\left(\frac{x_2-x_1}{-x_0}\right)g_\alpha^{a_2,a_1,a_3,a_4}(\tilde{w}_{(a_2)},\tilde{w}_{(a_1)},\tilde{w}_{(a_3)},\mathcal{B}(\mathcal{B}([\mathcal{Z}]_P));x_1,x_2)\\
&\quad = x_2^{-1}\delta\left(\frac{x_1-x_0}{x_2}\right)h_\alpha^{a_2,a_1,a_3,a_4}(\tilde{w}_{(a_2)},\tilde{w}_{(a_1)},\tilde{w}_{(a_3)},\mathcal{F}(\mathcal{B}([\mathcal{Z}]_P));x_0,x_2) \qquad (3.75)
\end{aligned}$$

for $\mathcal{Z}\in\coprod_{a_5\in\mathcal{A}}\mathcal{V}_{a_1a_5}^{a_4}\otimes\mathcal{V}_{a_2a_3}^{a_5}$ and $\alpha\in\mathbb{A}(a_2,a_1,a_3,a_4)$. Since $\mathcal{B}$ is isomorphic, by (3.64) we see that $\mathcal{B}([\mathcal{Z}]_P)$ in (3.71)-(3.75) can be any element in $\pi_P(\coprod_{a_5\in\mathcal{A}}\mathcal{V}_{a_2a_5}^{a_4}\otimes\mathcal{V}_{a_1a_3}^{a_5})$. So the Jacobi identity holds for the ordered triple $(\tilde{w}_{(a_2)},\tilde{w}_{(a_1)},\tilde{w}_{(a_3)})$. ∎

Theorem 3.7. *Assume that the assumptions of Theorem 3.1 hold, then the Jacobi identity holds for the ordered triple* $(\tilde{w}_{(a_1)},\tilde{w}_{(a_3)},\tilde{w}_{(a_2)})$.

Proof. Consider any $a_4\in\mathcal{A}$, $w'_{(a_4)}\in(W^{a_4})'$ and $\mathcal{Z}\in\coprod_{a_5\in\mathcal{A}}\mathcal{V}_{a_1a_5}^{a_4}\otimes\mathcal{V}_{a_2a_3}^{a_5}$. Observe that

$$e^{-x_2L(1)}w'_{(a_4)}\in(W^{a_4})'[x_2].$$

Then it can be easily derived that Proposition 3.3 holds with $w'_{(a_4)}$ replaced by $e^{-x_2L(1)}w'_{(a_4)}$. In particular, replacing $w'_{(a_4)}$ by $e^{-x_2L(1)}w'_{(a_4)}$ in (3.6), we get a multivalued analytic function

$$\Phi(e^{-x_2L(1)}w'_{(a_4)},\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)},[\mathcal{Z}]_P;z_1,z_2)\in\mathbb{G}^{a_1,a_2,a_3,a_4}. \qquad (3.76)$$

Moreover, replacing the complex variables (z_1,z_2) by $(z_1-z_2,-z_2)$ in (3.76), we get a multivalued analytic function on $M^2=\{(z_1,z_2)\in\mathbb{C}^2\mid z_1,z_2\neq 0, z_1\neq z_2\}$, which shall simply be denoted by

$$\Phi(e^{-x_2L(1)}w'_{(a_4)},\tilde{w}_{(a_1)},\tilde{w}_{(a_2)},\tilde{w}_{(a_3)},[\mathcal{Z}]_P;z_1-z_2,-z_2). \qquad (3.77)$$

Consider the simply connected region in $\mathbb{C}^2$ obtained by cutting the region $|z_1-z_2|>|z_2|>0$ along the intersection of this region with

$$\{(z_1,z_2)\in\mathbb{C}^2\mid z_2\in(-\infty,0]\}\cup\{(z_1,z_2)\in\mathbb{C}^2\mid z_1-z_2\in[0,+\infty)\}.$$

We denote it by R_5. Replacing $w'_{(a_4)}$ by $e^{-x_2L(1)}w'_{(a_4)}$, and then (z_1, z_2) by $(z_1 - z_2, -z_2)$ in (3.7), we see that

$$\langle e^{-x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)},x_2^n=e^{n\log(-z_2)}} \tag{3.78}$$

is a branch of $\Phi(e^{-x_2L(1)}w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1-z_2, -z_2)$ on the region R_5. Moreover, the skew-symmetry isomorphism implies

$$\begin{aligned}
&\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1},x_2^n=e^{n\log z_2}}\\
&\quad= \langle e^{x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, e^{-\pi i}x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)},x_2^n=e^{n\log z_2}}\\
&\quad= \langle e^{-x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)},x_2^n=e^{n\log(-z_2)}}
\end{aligned} \tag{3.79}$$

on the region $\{(z_1, z_2) \in \mathbb{C}^2 \mid \mathrm{Re}z_1 > -\mathrm{Re}z_2 > 0, \mathrm{Im}z_1 > -\mathrm{Im}z_2 > 0\}$. And observing that the single-valued analytic function

$$\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1},x_2^n=e^{n\log z_2}} \tag{3.80}$$

on the region $\{(z_1, z_2) \in \mathbb{C}^2 \mid \mathrm{Re}z_1 > -\mathrm{Re}z_2 > 0, \mathrm{Im}z_1 > -\mathrm{Im}z_2 > 0\}$ can be naturally analytically extended to the region R_1, we can conclude that the single-valued analytic function (3.80) on R_1 is a branch of $\Phi(e^{-x_2L(1)}w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1 - z_2, -z_2)$ on R_1.

Let (a_0, b_0), (a_1, b_1), (a_2, b_2) and (a_3, b_3) be four pairs of fixed positive real numbers satisfying

$$\begin{aligned}
&a_0 > b_0 > a_0 - b_0 > 0, \quad a_1 > a_1 - b_1 > b_1 > 0,\\
&b_2 > b_2 - a_2 > a_2 > 0, \quad b_3 > a_3 > b_3 - a_3 > 0.
\end{aligned} \tag{3.81}$$

Then we shall obtain branches of $\Phi(e^{-x_2L(1)}w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1 - z_2, -z_2)$ by analytical continuations along curves.

First of all, we consider the simply connected region

$$\begin{aligned}
\mathfrak{G}' \;=\; &\mathbb{C}^2\backslash\big(\{(z_1, z_2) \in \mathbb{C}^2 \mid z_1 \in [0, +\infty)\} \cup \{(z_1, z_2) \in \mathbb{C}^2 \mid z_2 \in [0, +\infty)\} \cup\\
&\{(z_1, z_2) \in \mathbb{C}^2 \mid z_1 - z_2 \in [0, +\infty)\}\big).
\end{aligned}$$

Define a path $\gamma : \ [0,1] \to \mathfrak{G}'$ by

$$
\begin{aligned}
\gamma(t) &= (\tilde{z}_1(t), \tilde{z}_2(t)) \\
&= \begin{cases}
\Big((a_0(1-7t)+7a_1t)e^{\frac{1}{4}\pi i},\ \ (b_0(1-7t)+7b_1t)e^{\frac{1}{4}\pi i}\Big) & t \in [0, \frac{1}{7}], \\
\Big(a_1 e^{\frac{1}{4}\pi i},\ \ b_1 e^{\frac{1}{4}\pi i + (7t-1)\pi i}\Big) & t \in (\frac{1}{7}, \frac{2}{7}], \\
\Big((a_1(3-7t)+a_2(7t-2))e^{\frac{1}{4}\pi i},\ \ (b_1(3-7t)+b_2(7t-2))e^{\frac{5}{4}\pi i}\Big) & t \in (\frac{2}{7}, \frac{3}{7}], \\
\Big(a_2 e^{\frac{1}{4}\pi i + (7t-3)\pi i},\ \ b_2 e^{\frac{5}{4}\pi i}\Big) & t \in (\frac{3}{7}, \frac{4}{7}], \\
\Big(b_2 e^{\frac{5}{4}\pi i} + (b_2 - a_2) e^{\frac{1}{4}\pi i + (7t-4)\pi i},\ \ b_2 e^{\frac{5}{4}\pi i}\Big) & t \in (\frac{4}{7}, \frac{5}{7}], \\
\Big((2b_2 - a_2) e^{\frac{5}{4}\pi i - (7t-5)\pi i},\ \ b_2 e^{\frac{5}{4}\pi i - (7t-5)\pi i}\Big) & t \in (\frac{5}{7}, \frac{6}{7}], \\
\Big((7(2b_2-a_2)(1-t) + a_0(7t-6))e^{\frac{1}{4}\pi i}, & \\
\qquad (7b_2(1-t) + b_0(7t-6))e^{\frac{1}{4}\pi i}\Big) & t \in (\frac{6}{7}, 1].
\end{cases}
\end{aligned}
\tag{3.82}
$$

See Figure 1 for an illustration. Then $\gamma(t) \subset \mathfrak{G}'$. We choose a simply connected region

$$D_t = \{(z_1, z_2) \in \mathbb{C}^2 \mid \max(|z_1 - \tilde{z}_1(t)|, |z_2 - \tilde{z}_2(t)|) < \varepsilon_t\} \tag{3.83}$$

for each $t \in [0,1]$, where ε_t is a sufficiently small positive real number for each $t \in [0,1]$ such that

$$D_0 \subset \mathfrak{G}' \cap \{(z_1, z_2) \in \mathbb{C}^2 \mid \mathrm{Re} z_1 > \mathrm{Re} z_2 > \mathrm{Re}(z_1 - z_2) > 0,\ \mathrm{Im} z_1 > \mathrm{Im} z_2 > \mathrm{Im}(z_1 - z_2) > 0\},$$

$$D_t \subset \mathfrak{G}' \cap \{(z_1, z_2) \in \mathbb{C}^2 \mid |z_1| > |z_2| > 0\} \quad \text{for } t \in (0, \frac{2}{7}),$$

$$D_{\frac{2}{7}} \subset \mathfrak{G}' \cap \{(z_1, z_2) \in \mathbb{C}^2 \mid \mathrm{Re} z_1 > -\mathrm{Re} z_2 > 0,\ \mathrm{Im} z_1 > -\mathrm{Im} z_2 > 0\},$$

$$D_t \subset \mathfrak{G}' \cap \{(z_1, z_2) \in \mathbb{C}^2 \mid \mathrm{Re} z_1 > 0 > \mathrm{Re} z_2,\ \mathrm{Im} z_1 > 0 > \mathrm{Im} z_2\} \quad \text{for } t \in (\frac{2}{7}, \frac{3}{7}),$$

$$D_{\frac{3}{7}} \subset \mathfrak{G}' \cap \{(z_1, z_2) \in \mathbb{C}^2 \mid -\mathrm{Re} z_2 > \mathrm{Re} z_1 > 0,\ -\mathrm{Im} z_2 > \mathrm{Im} z_1 > 0\},$$

$$D_t \subset \mathfrak{G}' \cap \{(z_1, z_2) \in \mathbb{C}^2 \mid |z_2| > |z_1| > 0\} \quad \text{for } t \in (\frac{3}{7}, \frac{4}{7}),$$

$$D_{\frac{4}{7}} \subset \mathfrak{G}' \cap \{(z_1, z_2) \in \mathbb{C}^2 \mid \mathrm{Re} z_2 < \mathrm{Re}(z_2 - z_1) < \mathrm{Re} z_1 < 0,\ \mathrm{Im} z_2 < \mathrm{Im}(z_2 - z_1) < \mathrm{Im} z_1 < 0\},$$

$$D_t \subset \mathfrak{G}' \cap \{(z_1, z_2) \in \mathbb{C}^2 \mid |z_2| > |z_1 - z_2| > 0\} \quad \text{for } t \in (\frac{4}{7}, 1),$$

$$D_1 = D_0.$$

With some straightforward calculations, the existence of ε_t can be easily verified. We omit the details here except that we shall write more about ε_t for $t \in (\frac{3}{7}, \frac{4}{7})$. Note that $|\tilde{z}_1(t)| = a_2$ and $|\tilde{z}_2(t)| = b_2$ for $t \in (\frac{3}{7}, \frac{4}{7})$. So for each $t \in (\frac{3}{7}, \frac{4}{7})$, to ensure that $D_t \subset \mathfrak{G}'$, we must have

Figure 1: $\gamma(t)$



$\varepsilon_t < a_2$, which further implies $\varepsilon_t < \frac{1}{2}b_2$ by (3.81). Thus $\mathrm{Re} z_2 < 0$ and $\mathrm{Im} z_2 < 0$ for any $(z_1, z_2) \in D_t$ with $t \in (\frac{3}{7}, \frac{4}{7})$. With these simply connected regions, we can see that

$$f_t = \langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}} \tag{3.84}$$

is a single-valued analytic function on the region D_t for each $t \in [0, \frac{2}{7}]$;

$$f_t = \langle e^{-x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)}, x_2^n=e^{n\log(-z_2)}} \tag{3.85}$$

is a single-valued analytic function on the region D_t for each $t \in (\frac{2}{7}, \frac{3}{7}]$;

$$f_t = \langle e^{-x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log(-z_2)}} \tag{3.86}$$

is a single-valued analytic function on the region D_t for each $t \in (\frac{3}{7}, \frac{4}{7}]$; and

$$f_t = \langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)}, x_2^n=e^{n\log z_2}} \tag{3.87}$$

is a single-valued analytic function on the region D_t for each $t \in (\frac{4}{7}, 1]$. Next, we shall show that $\{(f_t, D_t) : 0 \leq t \leq 1\}$ is an analytic continuation along γ.

Firstly, it can be derived from the skew-symmetry property that on the region $D_{\frac{2}{7}}$,

$$\begin{aligned}
f_{\frac{2}{7}} &= \langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}} \\
&= \langle e^{x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, e^{-\pi i}x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)}, x_2^n=e^{n\log z_2}} \\
&= \langle e^{-x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)}, x_2^n=e^{n\log(-z_2)}}.
\end{aligned} \tag{3.88}$$

Secondly, replacing $(w'_{(a_4)}, z_1, z_2)$ by $(e^{-x_2L(1)}w'_{(a_4)}, z_1 - z_2, -z_2)$ in (3.11), we can derive that

$$\begin{aligned}
f_{\frac{3}{7}} &= \langle e^{-x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)}, x_2^n=e^{n\log(-z_2)}} \\
&= \langle e^{-x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log(-z_2)}}
\end{aligned} \tag{3.89}$$

on the region $D_{\frac{3}{7}}$. Thirdly, we shall prove that on the region $D_{\frac{4}{7}}$,

$$\begin{aligned}
f_{\frac{4}{7}} &= \langle e^{-x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log(-z_2)}} \\
&= \langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)}, x_2^n=e^{n\log z_2}}.
\end{aligned} \tag{3.90}$$

By skew-symmetry, we have

$$\begin{aligned}
&\langle e^{-x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{I}}((\widetilde{\Omega^{-1}})^{(1)}\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_1, x_0)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log(-z_1)}, x_0^n=e^{n\log(z_1-z_2)}} \\
&\quad = \langle e^{-x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; e^{\pi i}x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log(-z_1)}, x_2^n=e^{n\log(-z_2)}} \\
&\quad = \langle e^{-x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log(-z_2)}}
\end{aligned} \tag{3.91}$$

on the region $D_{\frac{4}{7}}$. Moreover, note that $[\mathcal{Z}]_P \in \pi_P(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4}\otimes \mathcal{V}_{a_2a_3}^{a_5})$ implies

$$\mathcal{B}^{-1}(\mathcal{B}^{-1}([\mathcal{Z}]_P)) \in \pi_P(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4}\otimes \mathcal{V}_{a_2a_3}^{a_5}).$$

So replacing $[\mathcal{Z}]_P$ by $\mathcal{B}^{-1}(\mathcal{B}^{-1}([\mathcal{Z}]_P))$, $(w'_{(a_4)}, z_1, z_2)$ by $(e^{-x_2L(1)}w'_{(a_4)}, z_1-z_2, -z_2)$ in (3.12), we obtain that

$$\begin{aligned}
&\langle e^{-x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}^{-1}([\mathcal{Z}]_P)))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_2, x_0)\rangle_{W^{a_4}}\bigg|_{\substack{x_0^n=e^{n\log(z_1-z_2)}\\ x_2^n=e^{n\log(-z_2)}}}\\
&\quad= \langle e^{-x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{I}}((\widetilde{\Omega^{-1}})^{(1)}\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_1, x_0)\rangle_{W^{a_4}}\bigg|_{\substack{x_1^n=e^{n\log(-z_1)}\\ x_0^n=e^{n\log(z_1-z_2)}}}
\end{aligned}\tag{3.92}$$

on the region $D_{\frac{4}{7}}$. Furthermore, by the Moore-Seiberg equations (2.50) and (2.51), we have

$$\mathcal{F}\tilde{\Omega}^{(4)} = \tilde{\Omega}^{(2)}\mathcal{F}^{-1}\mathcal{F}(\widetilde{\Omega^{-1}})^{(3)}\mathcal{F}\tilde{\Omega}^{(4)} = \tilde{\Omega}^{(2)}\mathcal{F}^{-1}(\widetilde{\Omega^{-1}})^{(1)}\mathcal{F}(\widetilde{\Omega^{-1}})^{(4)}\tilde{\Omega}^{(4)} = \tilde{\Omega}^{(2)}\mathcal{B}^{-1}. \tag{3.93}$$

So this together with the skew-symmetry isomorphism implies

$$\begin{aligned}
&\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_0, x_2)\rangle_{W^{a_4}}\bigg|_{\substack{x_0^n=e^{n\log(z_1-z_2)}\\ x_2^n=e^{n\log z_2}}}\\
&\quad= \langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\tilde{\Omega}^{(2)}\mathcal{B}^{-1}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_0, x_2)\rangle_{W^{a_4}}\bigg|_{\substack{x_0^n=e^{n\log(z_1-z_2)}\\ x_2^n=e^{n\log z_2}}}\\
&\quad= \langle e^{x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}^{-1}([\mathcal{Z}]_P)))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; e^{-\pi i}x_2, x_0)\rangle_{W^{a_4}}\bigg|_{\substack{x_0^n=e^{n\log(z_1-z_2)}\\ x_2^n=e^{n\log z_2}}}\\
&\quad= \langle e^{-x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}^{-1}([\mathcal{Z}]_P)))(\tilde{w}_{(a_2)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}; x_2, x_0)\rangle_{W^{a_4}}\bigg|_{\substack{x_0^n=e^{n\log(z_1-z_2)}\\ x_2^n=e^{n\log(-z_2)}}}
\end{aligned}\tag{3.94}$$

on the region $D_{\frac{4}{7}}$. Therefore, (3.90) holds by (3.91), (3.92) and (3.94).

So to sum up, $\{(f_t, D_t) : 0 \le t \le 1\}$ is an analytic continuation along γ.

Since $\mathfrak{G}'$ is simply connected, $\gamma \subset \mathfrak{G}'$ and $\gamma(0) = \gamma(1)$, we can derive that $f_0 = f_1$ on the region $D_0 \cap D_1 = D_0$. Moreover, since f_0 and f_1 are both single-valued analytic functions on the domain S_1 which contains D_0, we deduce that $f_0 = f_1$ on S_1. Namely,

$$\begin{aligned}
&\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_1, x_2)\rangle_{W^{a_4}}|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}}\\
&\quad= \langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_0, x_2)\rangle_{W^{a_4}}|_{x_0^n=e^{n\log(z_1-z_2)}, x_2^n=e^{n\log z_2}}
\end{aligned}\tag{3.95}$$

on the region S_1. Furthermore, since the first line of (3.95) defined on R_1 is a branch of (3.77), and the second line of (3.95) on S_1 can be naturally analytically extended to the region R_3, we conclude that

$$\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_0, x_2)\rangle_{W^{a_4}}|_{x_0^n=e^{n\log(z_1-z_2)}, x_2^n=e^{n\log z_2}} \tag{3.96}$$

on R_3 is a branch of $\Phi(e^{-x_2L(1)}w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1 - z_2, -z_2)$ on R_3.

Then, by skew-symmetry isomorphism and by (3.95), we can deduce that on the region S_1,

$$\begin{aligned}
&\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\tilde{\Omega}^{(1)}\mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_3)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}; x_0, x_1)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_2-z_1)}, x_1^n=e^{n\log z_1}}\\
&\quad= \langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; e^{-\pi i}x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_2-z_1)}, x_2^n=e^{n\log z_2}}\\
&\quad= \langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)}, x_2^n=e^{n\log z_2}}\\
&\quad= \langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}}. \qquad (3.97)
\end{aligned}$$

Since the last line of (3.97) defined on R_1 is a branch of (3.77), and the first line of (3.97) on S_1 can be naturally analytically extended to the region R_4, we conclude that

$$\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\tilde{\Omega}^{(1)}\mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_3)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}; x_0, x_1)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_2-z_1)}, x_1^n=e^{n\log z_1}} \qquad (3.98)$$

on R_4 is a branch of $\Phi(e^{-x_2L(1)}w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1 - z_2, -z_2)$ on R_4.

Next, we consider the simply connected region

$$\begin{aligned}
\mathfrak{G}'' \;=\; &\mathbb{C}^2\backslash\big(\{(z_1, z_2) \in \mathbb{C}^2 \mid z_1 \in [0, +\infty)\} \cup \{(z_1, z_2) \in \mathbb{C}^2 \mid z_2 \in [0, +\infty)\}\\
&\cup\{(z_1, z_2) \in \mathbb{C}^2 \mid z_2 - z_1 \in [0, +\infty)\}\big).
\end{aligned}$$

Define a path $\sigma :\ [0, 1] \to \mathfrak{G}''$ by

$$\begin{aligned}
&\sigma(t) = (\tilde{z}_1(t), \tilde{z}_2(t))\\
&\quad= \begin{cases}
\Big(a_3e^{\frac{1}{4}\pi i},\ \ a_3e^{\frac{1}{4}\pi i} + (b_3 - a_3)e^{\frac{1}{4}\pi i+7t\pi i}\Big) & t \in [0, \frac{1}{7}],\\
\Big((a_3(2 - 7t) + a_0(7t - 1))e^{\frac{1}{4}\pi i}, & \\
\qquad ((2a_3 - b_3)(2 - 7t) + b_0(7t - 1))e^{\frac{1}{4}\pi i}\Big) & t \in (\frac{1}{7}, \frac{2}{7}],\\
\Big((a_0(3 - 7t) + a_1(7t - 2))e^{\frac{1}{4}\pi i},\ \ (b_0(3 - 7t) + b_1(7t - 2))e^{\frac{1}{4}\pi i}\Big) & t \in (\frac{2}{7}, \frac{3}{7}],\\
\Big(a_1e^{\frac{1}{4}\pi i},\ \ b_1e^{\frac{1}{4}\pi i+(7t-3)\pi i}\Big) & t \in (\frac{3}{7}, \frac{4}{7}],\\
\Big((a_1(5 - 7t) + a_2(7t - 4))e^{\frac{1}{4}\pi i},\ \ (b_1(5 - 7t) + b_2(7t - 4))e^{\frac{5}{4}\pi i}\Big) & t \in (\frac{4}{7}, \frac{5}{7}],\\
\Big(a_2e^{\frac{1}{4}\pi i},\ \ b_2e^{\frac{5}{4}\pi i-(7t-5)\pi i}\Big) & t \in (\frac{5}{7}, \frac{6}{7}],\\
\Big((7a_2(1 - t) + a_3(7t - 6))e^{\frac{1}{4}\pi i},\ \ (7b_2(1 - t) + b_3(7t - 6))e^{\frac{1}{4}\pi i}\Big) & t \in (\frac{6}{7}, 1].
\end{cases} \qquad (3.99)
\end{aligned}$$

See Figure 2 for an illustration. Then $\sigma(t) \subset \mathfrak{G}''$. We also choose a simply connected region

$$E_t = \{(z_1, z_2) \in \mathbb{C}^2 \mid \max(|z_1 - \tilde{z}_1(t)|, |z_2 - \tilde{z}_2(t)|) < \epsilon_t\}, \qquad (3.100)$$

for each $t \in [0, 1]$, where ϵ_t is a sufficiently small positive real number for each $t \in [0, 1]$ such that

$$E_0 \subset \mathfrak{G}'' \cap \{(z_1, z_2) \in \mathbb{C}^2 \mid \mathrm{Re} z_2 > \mathrm{Re} z_1 > \mathrm{Re}(z_2 - z_1) > 0,\ \mathrm{Im} z_2 > \mathrm{Im} z_1 > \mathrm{Im}(z_2 - z_1) > 0\},$$



Figure 2: $\sigma(t)$

$$\begin{gathered}
E_t \subset \mathfrak{G}'' \cap \{(z_1, z_2) \in \mathbb{C}^2 \mid |z_1| > |z_1 - z_2| > 0\} \quad \text{for } t \in (0, \frac{2}{7}),\\
E_{\frac{2}{7}} \subset \mathfrak{G}'' \cap S_1,\\
E_t \subset \mathfrak{G}'' \cap \{(z_1, z_2) \in \mathbb{C}^2 \mid |z_1| > |z_2| > 0\} \quad \text{for } t \in (\frac{2}{7}, \frac{4}{7}),\\
E_{\frac{4}{7}} \subset \mathfrak{G}'' \cap \{(z_1, z_2) \in \mathbb{C}^2 \mid \mathrm{Re} z_1 > -\mathrm{Re} z_2 > 0,\ \mathrm{Im} z_1 > -\mathrm{Im} z_2 > 0\},\\
E_t \subset \mathfrak{G}'' \cap \{(z_1, z_2) \in \mathbb{C}^2 \mid \mathrm{Re} z_1 > 0 > \mathrm{Re} z_2,\ \mathrm{Im} z_1 > 0 > \mathrm{Im} z_2\} \quad \text{for } t \in (\frac{4}{7}, \frac{5}{7}),\\
E_{\frac{5}{7}} \subset \mathfrak{G}'' \cap \{(z_1, z_2) \in \mathbb{C}^2 \mid -\mathrm{Re} z_2 > \mathrm{Re} z_1 > 0,\ -\mathrm{Im} z_2 > \mathrm{Im} z_1 > 0\},\\
E_t \subset \mathfrak{G}'' \cap \{(z_1, z_2) \in \mathbb{C}^2 \mid |z_2| > |z_1| > 0\} \quad \text{for } t \in (\frac{5}{7}, 1),\\
E_1 = E_0.
\end{gathered}$$

Thus

$$g_t = \langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\tilde{\Omega}^{(1)}\mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_3)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}; x_0, x_1)\rangle_{W^{a_4}}\Big|_{\substack{x_0^n = e^{n \log(z_2 - z_1)}\\ x_1^n = e^{n \log z_1}}} \tag{3.101}$$

is a single-valued analytic function on the region E_t for each $t \in [0, \frac{2}{7}]$;

$$g_t = \langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_1, x_2)\rangle_{W^{a_4}}|_{x_1^n = e^{n \log z_1}, x_2^n = e^{n \log z_2}} \tag{3.102}$$

is a single-valued analytic function on the region E_t for each $t \in (\frac{2}{7}, \frac{4}{7}]$;

$$g_t = \langle e^{-x_2 L(1)} w'_{(a_4)}, (\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}|_{x_0^n = e^{n \log(z_1 - z_2)}, x_2^n = e^{n \log(-z_2)}} \tag{3.103}$$

is a single-valued analytic function on the region E_t for each $t \in (\frac{4}{7}, \frac{5}{7}]$; and

$$g_t = \langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_3)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}; x_2, x_1)\rangle_{W^{a_4}}|_{x_1^n = e^{n \log z_1}, x_2^n = e^{n \log z_2}} \tag{3.104}$$

is a single-valued analytic function on the region E_t for each $t \in (\frac{5}{7}, 1]$. Next, we shall show that $\{(g_t, E_t) : 0 \leq t \leq 1\}$ is an analytic continuation along γ.

Firstly, by (3.97) we have

$$\begin{aligned}
g_{\frac{2}{7}} &= \langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\tilde{\Omega}^{(1)}\mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_3)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}; x_0, x_1)\rangle_{W^{a_4}}|_{x_0^n = e^{n \log(z_2 - z_1)}, x_1^n = e^{n \log z_1}}\\
&= \langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_1, x_2)\rangle_{W^{a_4}}|_{x_1^n = e^{n \log z_1}, x_2^n = e^{n \log z_2}} \qquad (3.105)
\end{aligned}$$

on the region $E_{\frac{2}{7}}$. Secondly, it can be derived from the skew-symmetry isomorphism that on the region $E_{\frac{4}{7}}$,

$$\begin{aligned}
g_{\frac{4}{7}} &= \langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_1, x_2)\rangle_{W^{a_4}}|_{x_1^n = e^{n \log z_1}, x_2^n = e^{n \log z_2}}\\
&= \langle e^{x_2 L(1)} w'_{(a_4)}, (\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, e^{-\pi i} x_2)\rangle_{W^{a_4}}|_{x_0^n = e^{n \log(z_1 - z_2)}, x_2^n = e^{n \log z_2}}\\
&= \langle e^{-x_2 L(1)} w'_{(a_4)}, (\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}|_{x_0^n = e^{n \log(z_1 - z_2)}, x_2^n = e^{n \log(-z_2)}}. \quad (3.106)
\end{aligned}$$

Thirdly, we shall prove that on the region $E_{\frac{5}{7}}$,

$$\begin{aligned} g_{\frac{5}{7}} &= \langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_3)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}; x_2, x_1)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}} \\ &= \langle e^{-x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)}, x_2^n=e^{n\log(-z_2)}}. \end{aligned} \tag{3.107}$$

Replacing $(w'_{(a_4)}, z_1, z_2)$ by $(e^{-x_2L(1)}w'_{(a_4)}, z_1-z_2, -z_2)$ in (3.11), we can derive that

$$\begin{aligned} &\langle e^{-x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{P}}([\mathcal{Z}]_P))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)}, x_2^n=e^{n\log(-z_2)}} \\ &\quad = \langle e^{-x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log(-z_2)}} \end{aligned} \tag{3.108}$$

on the region $E_{\frac{5}{7}}$. Moreover, by the Moore-Seiberg equations (2.50) and (2.51), we have

$$\mathcal{B}\tilde{\Omega}^{(4)} = \mathcal{F}^{-1}\tilde{\Omega}^{(1)}\mathcal{F}\tilde{\Omega}^{(4)} = \tilde{\Omega}^{(3)}\mathcal{F}. \tag{3.109}$$

So by skew-symmetry we have

$$\begin{aligned} &\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_3)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}; x_2, x_1)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}} \\ &\quad = \langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\tilde{\Omega}^{(3)}\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_3)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}; x_2, x_1)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}} \\ &\quad = \langle e^{x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_1, e^{-\pi i}x_2)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}} \\ &\quad = \langle e^{-x_2L(1)}w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log(-z_2)}} \end{aligned} \tag{3.110}$$

on the region $E_{\frac{5}{7}}$. Therefore, (3.107) holds by (3.108) and (3.110).

So to sum up, $\{(g_t, E_t) : 0 \leq t \leq 1\}$ is an analytic continuation along σ.

Since $\mathfrak{G}''$ is simply connected, $\sigma \subset \mathfrak{G}''$ and $\sigma(0) = \sigma(1)$, we can derive that $g_0 = g_1$ on the region $E_0 \cap E_1 = E_0$. Moreover, since g_0 and g_1 are both single-valued analytic functions on the domain S_2 which contains E_0, we deduce that $g_0 = g_1$ on S_2. Namely,

$$\begin{aligned} &\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\tilde{\Omega}^{(1)}\mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_3)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}; x_0, x_1)\rangle_{W^{a_4}}\Big|_{x_0^n=e^{n\log(z_2-z_1)}, x_1^n=e^{n\log z_1}} \\ &\quad = \langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_3)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}; x_2, x_1)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}} \end{aligned} \tag{3.111}$$

on the region S_2. Furthermore, since the first line of (3.111) defined on R_4 is a branch of (3.77), and the second line of (3.111) on S_2 can be naturally analytically extended to the region R_2, we conclude that

$$\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_3)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}; x_2, x_1)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}} \tag{3.112}$$

on R_2 is a branch of $\Phi(e^{-x_2L(1)}w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1-z_2, -z_2)$ on R_2.

In conclusion of (3.77)-(3.112), we see that the multivalued analytic functions

$$\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_1, x_2)\rangle_{W^{a_4}}\big|_{x_1=z_1, x_2=z_2}, \tag{3.113}$$

$$\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_3)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}; x_2, x_1)\rangle_{W^{a_4}}\big|_{x_1=z_1, x_2=z_2} \tag{3.114}$$

and

$$\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0=z_1-z_2, x_2=z_2} \tag{3.115}$$

are restrictions of $\Phi(e^{-x_2L(1)}w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1-z_2, -z_2)$ to their domains $|z_1| > |z_2| > 0$, $|z_2| > |z_1| > 0$ and $|z_2| > |z_1 - z_2| > 0$ respectively.

So choosing (3.80) as the preferred branch of $\Phi(e^{-x_2L(1)}w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1 - z_2, -z_2)$ on R_1, we see that $\Phi(e^{-x_2L(1)}w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1 - z_2, -z_2)$ is an element of $\mathbb{G}^{a_1,a_3,a_2,a_4}$. Moreover, by (3.95), (3.96), (3.97), (3.98), (3.111), (3.112), and by the definition of the preferred branches of an element of $\mathbb{G}^{a_1,a_3,a_2,a_4}$ on R_2 and R_3, we see that

$$\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_3)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}; x_2, x_1)\rangle_{W^{a_4}}\big|_{x_1^n=e^{n\log z_1}, x_2^n=e^{n\log z_2}} \tag{3.116}$$

and

$$\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_0, x_2)\rangle_{W^{a_4}}\big|_{x_0^n=e^{n\log(z_1-z_2)}, x_2^n=e^{n\log z_2}} \tag{3.117}$$

are the preferred branches of $\Phi(e^{-x_2L(1)}w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1 - z_2, -z_2)$ on R_2 and R_3 respectively.

Therefore, we have

$$\begin{aligned}\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_1, x_2)\rangle_{W^{a_4}}&\\ = \iota_{12}\Phi(e^{-x_2L(1)}w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1 - z_2, -z_2),&\end{aligned} \tag{3.118}$$

$$\begin{aligned}\langle w'_{(a_4)}, (\tilde{\mathbf{P}}(\mathcal{B}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_3)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}; x_2, x_1)\rangle_{W^{a_4}}&\\ = \iota_{21}\Phi(e^{-x_2L(1)}w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1 - z_2, -z_2),&\end{aligned} \tag{3.119}$$

$$\begin{aligned}\langle w'_{(a_4)}, (\tilde{\mathbf{I}}(\mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_0, x_2)\rangle_{W^{a_4}}&\\ = \iota_{20}\Phi(e^{-x_2L(1)}w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1 - z_2, -z_2).&\end{aligned} \tag{3.120}$$

Let $\{e_\alpha^{a_1,a_3,a_2,a_4}\}_{\alpha\in\mathbb{A}(a_1,a_3,a_2,a_4)}$ be a basis of $\mathbb{G}^{a_1,a_3,a_2,a_4}$ over the ring

$$\mathbb{C}[x_1, x_1^{-1}, x_2, x_2^{-1}, (x_1 - x_2)^{-1}]. \tag{3.121}$$

Then there exist unique

$$\begin{aligned}H_\alpha(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}, \tilde{\Omega}^{(4)}([\mathcal{Z}]_P); x_1, x_2)&\\ \in \mathbb{C}[x_1, x_1^{-1}, x_2, x_2^{-1}, (x_1 - x_2)^{-1}]&\end{aligned} \tag{3.122}$$

for $\alpha \in \mathbb{A}(a_1, a_3, a_2, a_4)$, such that only finitely many of them are nonzero and

$$\begin{aligned}&\Phi(e^{-x_2L(1)}w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}, \tilde{w}_{(a_3)}, [\mathcal{Z}]_P; z_1 - z_2, -z_2)\\&\quad = \sum_{\alpha\in\mathbb{A}(a_1,a_3,a_2,a_4)} H_\alpha(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}, \tilde{\Omega}^{(4)}([\mathcal{Z}]_P); x_1, x_2)e_\alpha^{a_1,a_3,a_2,a_4}.\end{aligned} \tag{3.123}$$

Recall that $\tilde{\Omega}^{(4)}$ is an isomorphism and that

$$\tilde{\Omega}^{(4)}\left(\pi_P(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5})\right) = \pi_P(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_3a_2}^{a_5}) \tag{3.124}$$

for any $a_1, \cdots, a_4 \in \mathcal{A}$. So we can define linear maps

$$\begin{aligned}&f_\alpha^{a_1,a_3,a_2,a_4}(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}) :\\&\qquad \pi_P(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_3a_2}^{a_5}) \to W^{a_4}[x_1, x_1^{-1}, x_2, x_2^{-1}][[x_2/x_1]],\end{aligned} \tag{3.125}$$

$$\begin{aligned}&g_\alpha^{a_1,a_3,a_2,a_4}(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}) :\\&\qquad \pi_P(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_3a_5}^{a_4} \otimes \mathcal{V}_{a_1a_2}^{a_5}) \to W^{a_4}[x_1, x_1^{-1}, x_2, x_2^{-1}][[x_1/x_2]],\end{aligned} \tag{3.126}$$

$$\begin{aligned}&h_\alpha^{a_1,a_3,a_2,a_4}(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}) :\\&\qquad \pi_I(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_3}^{a_5} \otimes \mathcal{V}_{a_5a_2}^{a_4}) \to W^{a_4}[x_0, x_0^{-1}, x_2, x_2^{-1}][[x_0/x_2]]\end{aligned} \tag{3.127}$$

by

$$\begin{aligned}&\langle w'_{(a_4)}, f_\alpha^{a_1,a_3,a_2,a_4}(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}, \tilde{\Omega}^{(4)}([\mathcal{Z}]_P); x_1, x_2)\rangle_{W^{a_4}}\\&\quad = \iota_{12}H_\alpha(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}, \tilde{\Omega}^{(4)}([\mathcal{Z}]_P); x_1, x_2),\end{aligned} \tag{3.128}$$

$$\begin{aligned}&\langle w'_{(a_4)}, g_\alpha^{a_1,a_3,a_2,a_4}(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}, \mathcal{B}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)); x_1, x_2)\rangle_{W^{a_4}}\\&\quad = \iota_{21}H_\alpha(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}, \tilde{\Omega}^{(4)}([\mathcal{Z}]_P); x_1, x_2),\end{aligned} \tag{3.129}$$

$$\begin{aligned}&\langle w'_{(a_4)}, h_\alpha^{a_1,a_3,a_2,a_4}(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}, \mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)); x_0, x_2)\rangle_{W^{a_4}}\\&\quad = \iota_{20}H_\alpha(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}, \tilde{\Omega}^{(4)}([\mathcal{Z}]_P); x_2 + x_0, x_2)\end{aligned} \tag{3.130}$$

for $w'_{(a_4)} \in (W^{a_4})'$, $\mathcal{Z} \in \coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5}$ and $\alpha \in \mathbb{A}(a_1, a_3, a_2, a_4)$. Then by (3.118), (3.119), (3.120) and (3.123), we have

$$\begin{aligned}&(\tilde{\mathbf{P}}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_1, x_2)\\&\quad = \sum_{\alpha\in\mathbb{A}(a_1,a_3,a_2,a_4)} f_\alpha^{a_1,a_3,a_2,a_4}(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}, \tilde{\Omega}^{(4)}([\mathcal{Z}]_P); x_1, x_2)\iota_{12}(e_\alpha^{a_1,a_3,a_2,a_4}),\end{aligned} \tag{3.131}$$

$$\begin{aligned}&(\tilde{\mathbf{P}}(\mathcal{B}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_3)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_2)}; x_2, x_1)\\&\quad = \sum_{\alpha\in\mathbb{A}(a_1,a_3,a_2,a_4)} g_\alpha^{a_1,a_3,a_2,a_4}(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}, \mathcal{B}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)); x_1, x_2)\iota_{21}(e_\alpha^{a_1,a_3,a_2,a_4}), \qquad (3.132)\end{aligned}$$

$$\begin{aligned}&(\tilde{\mathbf{I}}(\mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P))))(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}; x_0, x_2)\\&\quad = \sum_{\alpha\in\mathbb{A}(a_1,a_3,a_2,a_4)} h_\alpha^{a_1,a_3,a_2,a_4}(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}, \mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)); x_0, x_2)\iota_{20}(e_\alpha^{a_1,a_3,a_2,a_4}). \qquad (3.133)\end{aligned}$$

Moreover, by (3.122) and Proposition 2.1, we have

$$\begin{aligned}&x_0^{-1}\delta\left(\frac{x_1-x_2}{x_0}\right)\iota_{12}H_\alpha(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}, \tilde{\Omega}^{(4)}([\mathcal{Z}]_P); x_1, x_2)\\&\quad -x_0^{-1}\delta\left(\frac{x_2-x_1}{-x_0}\right)\iota_{21}H_\alpha(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}, \tilde{\Omega}^{(4)}([\mathcal{Z}]_P); x_1, x_2)\\&\quad = x_2^{-1}\delta\left(\frac{x_1-x_0}{x_2}\right)\iota_{20}H_\alpha(w'_{(a_4)}, \tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}, \tilde{\Omega}^{(4)}([\mathcal{Z}]_P); x_2+x_0, x_2) \qquad (3.134)\end{aligned}$$

for $\alpha \in \mathbb{A}(a_1, a_3, a_2, a_4)$. Since $w'_{(a_4)} \in (W^{a_4})'$ is arbitrary, by (3.128), (3.129), (3.130) and (3.134), the Jacobi identity holds for the ordered triple $(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)})$:

$$\begin{aligned}&x_0^{-1}\delta\left(\frac{x_1-x_2}{x_0}\right)f_\alpha^{a_1,a_3,a_2,a_4}(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}, \tilde{\Omega}^{(4)}([\mathcal{Z}]_P); x_1, x_2)\\&\quad -x_0^{-1}\delta\left(\frac{x_2-x_1}{-x_0}\right)g_\alpha^{a_1,a_3,a_2,a_4}(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}, \mathcal{B}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)); x_1, x_2)\\&\quad = x_2^{-1}\delta\left(\frac{x_1-x_0}{x_2}\right)h_\alpha^{a_1,a_3,a_2,a_4}(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)}, \mathcal{F}(\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)); x_0, x_2) \qquad (3.135)\end{aligned}$$

for $\mathcal{Z} \in \coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_2a_3}^{a_5}$ and $\alpha \in \mathbb{A}(a_1, a_3, a_2, a_4)$. By (3.124) we see that $\tilde{\Omega}^{(4)}([\mathcal{Z}]_P)$ in (3.131)-(3.135) can be any element in $\pi_P(\coprod_{a_5\in\mathcal{A}} \mathcal{V}_{a_1a_5}^{a_4} \otimes \mathcal{V}_{a_3a_2}^{a_5})$. So the Jacobi identity holds for the ordered triple $(\tilde{w}_{(a_1)}, \tilde{w}_{(a_3)}, \tilde{w}_{(a_2)})$. ∎

Proof of Theorem 3.1. Since the permutations (1 2) and (2 3) generate the symmetric group S_3, in summary of Theorems 3.6 and 3.7, we can conclude that the Jacobi identity holds for the triple $(\tilde{w}_{(a_{\tau(1)})}, \tilde{w}_{(a_{\tau(2)})}, \tilde{w}_{(a_{\tau(3)})})$ for any $\tau \in S_3$. Thus Theorem 3.1 holds. ∎

References

[C] L. Chen, On axiomatic approaches to intertwining operator algebras, to appear in *Communications in Contemporary Mathematics*, 1550051, 2015, DOI: 10.1142/S0219199715500510.

[DL1] C. Dong and J. Lepowsky, Abelian intertwining algebras–a generalization of vertex operator algebras, in: *Proc. AMS Summer Research Institute on Algebraic Groups and Their Generalizations, Pennsylvania State University*, 1991, Proc. Symp. Pure Math., Vol. 56, Part 2 (W. J. Haboush and B. J. Parshall, eds.). American Mathematical Society, Providence, 1994, pp. 261-293.

[DL2] C. Dong and J. Lepowsky, Generalized Vertex Algebras and Relative Vertex Operators, *Progress in Math.*, Vol. 112. Birkhäuser, Boston, 1993.

[FHL] I. B. Frenkel, Y. Z. Huang and J. Lepowsky, On axiomatic approaches to vertex operator algebras and modules, preprint 1989, *Memoirs. Amer. Math. Soc.* **104** (1993).

[FLM] I. B. Frenkel, J. Lepowsky and A. Meurman, Vertex operator algebras and the Monster, Pure and Appl. Math., Vol. 134, Academic Press, New York, 1988.

[G] H. Guo, On abelian intertwining algebras and their modules, Ph.D. Thesis, Rutgers University, 1995.

[H1] Y.-Z. Huang, Intertwining operator algebras, genus-zero modular functors and genus-zero conformal field theories, in: Operads: Proceedings of Renaissance Conferences, Contemporary Math., Vol. 202 (J. L. Loday, J. Stasheff, and A. A. Voronov, eds.); *Amer. Math. Soc., Provindence* 1997, pp. 335-355.

[H2] Y.-Z. Huang, Generalized rationality and a "Jacobi identity" for intertwining operator algebras, *Selecta Mathematica, New series* **6** (2000), 225-267.

[HL] Y.-Z. Huang and J. Lepowsky, A theory of tensor products for module categories for a vertex operator algebra, II, *Selecta Mathematica, New series* **1** (1995), 757-786.

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, CHINA
E-mail address: chenling2013@ucas.ac.cn